

## FIBRINOGEN RECEPTOR ANTAGONISTS

Patent Number: WO9725323

Publication date: 1997-07-17

Inventor(s): ALI FADIA EL-FEHAIL (US); SAMANEN JAMES (US); BONDINELL WILLIAM EDWARD (US); CALVO RAUL ROLANDO (US); KU THOMAS WEN-FU (US); MILLER WILLIAM HENRY (US); VENSILAVSKY JOSEPH WALTER (US); YELLIN TOBIAS OREGON (US)

Applicant(s): ALI FADIA EL FEHAIL (US); SMITHKLINE BEECHAM CORP (US); SAMANEN JAMES (US); BONDINELL WILLIAM EDWARD (US); CALVO RAUL ROLANDO (US); KU THOMAS WEN FU (US); MILLER WILLIAM HENRY (US); VENSILAVSKY JOSEPH WALTER (US); YELLIN TOBIAS OREGON (US)

Requested Patent: WO9725323

Application Number: WO1997US00326 19970109

Priority Number(s): US19960584030 19960111

IPC Classification: C07D401/12; C07D403/12; C07D295/135; A61K31/495; A61K31/445

EC Classification: C07D211/58; C07D211/62; C07D295/12A1; C07D295/20D1D

Equivalents: EP0885213 (WO9725323), A4, JP2001508757T

Cited Documents: DE4446301; DE4241632; WO9620173

### Abstract

This invention relates to compounds of formula (I), which are effective for inhibiting platelet aggregation, pharmaceutical compositions for effecting such activity, and a method for inhibiting platelet aggregation.

Data supplied from the esp@cenet database - I2

### Description

#### FIBRINOGEN RECEPTOR ANTAGONISTS

This invention relates to novel compounds which inhibit platelet aggregation, pharmaceutical compositions containing the compounds and methods of using the compounds.

#### Background of the Invention

Platelet aggregation is believed to be mediated primarily through the fibrinogen receptor, or GPIIb-IIIa platelet receptor complex, which is a member of a family of adhesion receptors referred to as integrins. It has been found that frequently the natural ligands of integrin receptors are proteins which contain an Arg-Gly-Asp sequence. Von Willebrand factor and fibrinogen, which are considered to be natural ligands for the GPIIb-IIIa receptor, possess an Arg-Gly-Asp (RGD in single letter amino acid code) sequence in their primary structure.

Functionally, these proteins are able to bind and crosslink GPIIb-IIIa receptors on adjacent platelets and thereby effect aggregation of platelets.

Fibronectin, vitronectin and thrombospondin are RGD-containing proteins which have also been demonstrated to bind to GPIIb-IIIa. Fibronectin is found in plasma and as a structural protein in the intracellular matrix. Binding between the structural proteins and GPITb-IIIa may function to cause

**BEST AVAILABLE COPY**


# PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6: C07D 401/12, 403/12, 295/135, A61K 31/495, 31/445		(11) International Publication Number: <b>WO 97/25323</b>	
		(43) International Publication Date: 17 July 1997 (17.07.97)	
(21) International Application Number: PCT/US97/00326		Henry (US/US): 333 Fell Lane, Schwenksville, PA 19473 (US); SAMANEN, James (US/US); Jug, Hollow Road, Phoenixville, PA 19460 (US); VENSILAVSKY, Joseph, Walter (US/US); 481 Brookwood Road, Wayne, PA 19087 (US); YELLEN, Tobias, Oregon (US/US); 517 Oriole Lane, Villanova, PA 19083 (US).	
(22) International Filing Date: 9 January 1997 (09.01.97)		(74) Agent: MCCARTHY, Mary, E. et al.; SmithKline Beecham Corporation, Corporate Intellectual Property, UW7220, 709 Swedeland Road, P.O. Box 1539, King of Prussia, PA 19406-0939 (US).	
(30) Priority Data: 08/584,030 11 January 1996 (11.01.96) US		(81) Designated States: JP, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).	
(60) Parent Application or Grant (63) Related by Continuation US Filed on 08/584,030 (CON) 11 January 1996 (11.01.96)		Published With international search report.	
(71) Applicant (for all designated States except US): SMITHKLINE BEECHAM CORPORATION (US/US); Corporate Intellectual Property, UW7220, 709 Swedeland Road, P.O. Box 1539, King of Prussia, PA 19406-0939 (US).			
(72) Inventors; and (73) Inventor Applicants (for US only): ALL, Roda, El-Schell (US/US); 5 North Green Acres Drive, Cherry Hill, NJ 08003 (US); BONDURIELLO, William, Edward (US/US); 512 Franklin Lane, Wayne, PA 19087 (US); CALVO, Raul, Rolando (US/US); 72 Orchard Court, Noyesford, PA 19068 (US); KU, Thomas, Wen-Pa (US/US); 1413 Southwind Way, Dresher, PA 19025 (US); MILLER, William,			

(54) Title: FIBRINOGEN RECEPTOR ANTAGONISTS



(57) Abstract

This invention relates to compounds of formula (I), which are effective for inhibiting platelet aggregation, pharmaceutical compositions for effecting such activity, and a method for inhibiting platelet aggregation.

### FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AN	Armenia	GB	United Kingdom	AW	Malawi
AT	Austria	GE	Georgia	AX	Mexico
AU	Australia	GR	Greece	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BO	Bolivia	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KC	Kazakhstan	RU	Russia Federation
CA	Canada	KG	Kyrgyzstan	SD	Sudan
CH	Switzerland	KR	Republic of Korea	SE	Sweden
CN	China	KR	Republic of Korea	SG	Singapore
CO	Colombia	KZ	Kazakhstan	SI	Slovenia
CZ	Czech Republic	LA	Laos	SK	Slovakia
DE	Germany	LB	Lebanon	SM	San Marino
DK	Denmark	LV	Latvia	SN	Senegal
EE	Estonia	LU	Luxembourg	SV	Sweden
ES	Spain	LT	Lithuania	TD	Togo
FI	Finland	LV	Latvia	TG	Togo
FR	France	MD	Republic of Moldova	TJ	Tajikistan
GA	Gabon	MG	Madagascar	TT	Trinidad and Tobago
		ML	Mali	UA	Ukraine
		MN	Mongolia	UG	Uganda
		MR	Mauritania	US	United States of America
				UZ	Uzbekistan
				VN	Viet Nam

## FIBRINOGEN RECEPTOR ANTAGONISTS

This invention relates to novel compounds which inhibit platelet aggregation, pharmaceutical compositions containing the compounds and methods of using the compounds.

### Background of the Invention

Platelet aggregation is believed to be mediated primarily through the fibrinogen receptor, or GPIIb-IIIa platelet receptor complex, which is a member of a family of adhesion receptors referred to as Integrins. It has been found that frequently the natural ligands of integrin receptors are proteins which contain an Arg-Gly-Asp sequence. Von Willebrand factor and fibrinogen, which are considered to be natural ligands for the GPIIb-IIIa receptor, possess an Arg-Gly-Asp (RGD in single letter amino acid code) sequence in their primary structure. Functionally, these proteins are able to bind and crosslink GPIIb-IIIa receptors on adjacent platelets and thereby effect aggregation of platelets.

Fibrinectin, vitronectin and thrombospondin are RGD-containing proteins which have also been demonstrated to bind to GPIIb-IIIa. Fibrinectin is found in plasma and as a structural protein in the intracellular matrix. Binding between the structural proteins and GPIIb-IIIa may function to cause platelets to adhere to damaged vessel walls.

Linear and cyclic peptides which bind to vitronectin and contain an RGD sequence are disclosed in WO 89/05150 (PCT US88/04403). EP 0 275 748 discloses linear tetra- to hexapeptides and cyclic hexa- to octapeptides which bind to the GPIIb-IIIa receptor and inhibit platelet aggregation. Other linear and cyclic peptides, the disclosure of which are incorporated herein by reference, are reported in EP-A 0 341 915. However, the peptide like structures of such inhibitors often pose problems, such as in drug delivery, metabolic stability and selectivity.

Inhibitors of the fibrinogen receptor which are not constructed of natural amino acid sequences are disclosed in EP-A 0 372,486, EP-A 0 381 033 and EP-A 0 478 363. WO 92/07568 (PCT/US91/08166) discloses fibrinogen receptor antagonists which mimic a conformational  $\gamma$ -turn in the RGD sequence by forming a monocyclic seven-membered ring structure. There remains a need, however, for novel

fibrinogen receptor antagonists (e.g., inhibitors of the GPIIb-IIIa protein) which have potent *in vivo* and *in vitro* effects and lack the peptide backbone structure of amino acid sequences.

The present invention discloses novel compounds. These compounds inhibit the GPIIb-IIIa receptor and inhibit platelet aggregation.

### Summary of the Invention

In one aspect this invention is a compound as described hereinafter in formula (I).

This invention is also a pharmaceutical composition for inhibiting platelet aggregation or clot formation, which comprises a compound of formula (I) and a pharmaceutically acceptable carrier.

This invention is further a method for inhibiting platelet aggregation in a mammal in need thereof, which comprises internally administering an effective amount of a compound of formula (I).


In another aspect, this invention provides a method for inhibiting reocclusion of an artery or vein in a mammal following fibrinolytic therapy, which comprises internally administering an effective amount of a fibrinolytic agent and a compound of formula (I). This invention is also a method for treating stroke, transient ischemia attacks, or myocardial infarction.

### Detailed Description of the Invention

This invention discloses compounds which inhibit platelet aggregation. The compounds of the instant invention are believed to interact favorably with the GPIIb-IIIa receptor.

Although not intending to be bound to any specific mechanism of action, these compounds are believed to inhibit the binding of fibrinogen to the platelet-bound fibrinogen receptor GPIIb-IIIa, and may interact with other adhesion proteins via antagonism of a putative RGD binding site.

The compounds of this invention are compounds of formula (I):




2


(I)

wherein:

R<sup>1</sup> is hydrogen, C<sub>1-6</sub>alkyl, HC≡C-CH<sub>2</sub>-, F<sub>3</sub>C-CH<sub>2</sub>-,



5



A<sup>1</sup>, A<sup>2</sup>, A<sup>3</sup>, A<sup>4</sup>, and A<sup>5</sup> independently are N or CH, with the proviso that A<sup>1</sup> and A<sup>4</sup> are not simultaneously N;

Y and Y<sup>1</sup> independently are hydrogen, C<sub>1-6</sub>alkyl, halo, CF<sub>3</sub>,

-(CH<sub>2</sub>)<sub>n</sub>-phenyl or together form a phenyl ring;

X is C(O)NR<sup>2</sup>, NR<sup>2</sup>C(O), SO<sub>2</sub>NR<sup>2</sup>, NR<sup>2</sup>SO<sub>2</sub>, C(O), C(R<sup>2</sup>)<sub>2</sub>, S, O, NR<sup>2</sup>, CR<sup>2</sup>OR<sup>2</sup>, C(O)C(R<sup>2</sup>)<sub>2</sub>, C(R<sup>2</sup>)<sub>2</sub>C(O), CH<sub>2</sub>CH<sub>2</sub>, CHOCH<sub>2</sub>, CH<sub>2</sub>CHOH, or CR<sup>2</sup>=CR<sup>2</sup>;

Q is a single bond or C(O);

Z is hydrogen, C<sub>1-4</sub>alkyl, OR<sup>2</sup>, CH<sub>2</sub>OR<sup>2</sup>, CH<sub>2</sub>CO<sub>2</sub>R<sup>2</sup>, CH<sub>2</sub>CONR<sup>2</sup>R<sup>2</sup>, CO<sub>2</sub>R<sup>2</sup>, CONR<sup>2</sup>R<sup>2</sup>, -CH=CH-, -C≡C-, COR<sup>2</sup>, or -(CH<sub>2</sub>)<sub>n</sub>-phenyl;

each n independently is 0-3; and

each R<sup>2</sup> independently is H or C<sub>1-6</sub>alkyl;


or a pharmaceutically acceptable salt thereof.

Also included in this invention are pharmaceutically acceptable addition salts, complexes or prodrugs of the compounds of this invention. Prodrugs are considered to be any covalently bonded carriers which release the active parent drug according to formula (I) *in vivo*.

In cases wherein the compounds of this invention may have one or more chiral centers, unless specified, this invention includes each unique nonracemic compound which may be synthesized and resolved by conventional techniques. In cases in which the compounds have unsaturated double bonds, both the *cis* (Z) and *trans* (E) are within the scope of this invention. The meaning of any substituent at

any one occurrence is independent of its meaning, or any other substituent's meaning, at any other occurrence, unless specified otherwise.

With reference to formula (II), compounds of formula (II) are preferred:



5

(II)

wherein:

Y and Y<sup>1</sup> independently are hydrogen, C<sub>1-6</sub>alkyl, halo, CF<sub>3</sub> or together form a phenyl ring;

X is C(O)NR<sup>2</sup>, NR<sup>2</sup>C(O), SO<sub>2</sub>NR<sup>2</sup>, NR<sup>2</sup>SO<sub>2</sub>, C(O), C(R<sup>2</sup>)<sub>2</sub>, S, O, or NR<sup>2</sup>;

Z is hydrogen, C<sub>1-4</sub>alkyl, OR<sup>2</sup>, CH<sub>2</sub>OR<sup>2</sup>, CH<sub>2</sub>CO<sub>2</sub>R<sup>2</sup>, or CO<sub>2</sub>R<sup>2</sup>; and

n is 0 or 1.

Preferably, the compounds of formula (II) are those wherein A<sup>1</sup> is N, A<sup>2</sup> is CH, X is C(O)NR<sup>2</sup> or NR<sup>2</sup>C(O), Z is hydrogen, R<sup>1</sup> is hydrogen or C<sub>1-4</sub>alkyl, and R<sup>2</sup> is hydrogen or C<sub>1-4</sub>alkyl.


A preferred compound of this invention is 4-[[4-(1-piperazinyl)phenyl]-amino]carbonyl-1-piperidineacetic acid or pharmaceutically acceptable salts thereof or C<sub>1-6</sub>alkyl esters thereof.

In the above description, C<sub>1-4</sub>alkyl is meant to include methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl and t-butyl. C<sub>1-6</sub>alkyl additionally includes pentyl, n-pentyl, isopentyl, neopentyl and hexyl and the simple aliphatic isomers thereof.

Certain radical groups are abbreviated herein. t-Bu refers to the tertiary butyl radical, Boc refers to the t-butyloxycarbonyl radical, Fmoc refers to the fluorenylmethoxycarbonyl radical, Ph refers to the phenyl radical, Cbz refers to the benzyloxycarbonyl radical, BrZ refers to the o-bromobenzyloxycarbonyl radical, ClZ refers to the o-chlorobenzyloxycarbonyl radical, Bzl refers to the benzyl radical, 4-MBzl refers to the 4-methyl benzyl radical, Me refers to methyl, Et refers to ethyl, Ac refers to acetyl, Alk refers to C<sub>1-6</sub>alkyl, Nph refers to 1- or 2-naphthyl and cHex refers to cyclohexyl. MeArg is N<sup>α</sup>-methyl arginine. Tet refers to 5-tetrazolyl.

Certain reagents are abbreviated herein. DCC refers to dicyclohexylcarbodiimide, DMAP refers to dimethylaminopyridine, DIEA refers to diisopropylethyl amine, EDC refers to N-ethyl-N'-(dimethylaminopropyl)-carbodiimide. HOBt refers to 1-hydroxybenzotriazole, THF refers to tetrahydrofuran, DMF refers to dimethyl formamide, NBS refers to N-bromosuccinimide, Pd/C refers to a palladium on carbon catalyst, PPA refers to l-propanephosphonic acid cyclic anhydride, DPPA refers to diphenylphosphoryl azide, BOP refers to benzotriazol-1-yloxy-tris(dimethylamino)phosphonium hexafluorophosphate, HF refers to hydrofluoric acid, TEA refers to triethylamine, TFA refers to trifluoroacetic acid, PCC refers to pyridinium chlorochromate.

The compounds of formula (I) are generally prepared by reacting a compound of the formula (III) with a compound of the formula (IV):



wherein A<sup>1</sup>, A<sup>2</sup>, A<sup>3</sup>, D, R<sup>1</sup>, R<sup>2</sup>, Q, Z, and n are as defined in formula (I), with any reactive functional groups protected; and

L<sup>1</sup> and L<sup>2</sup> are functional groups which are capable of reacting to form the X linkage;

and thereafter removing any protecting groups, and optionally forming a pharmaceutically acceptable salt.

It will be apparent that the precise identity of L<sup>1</sup> and L<sup>2</sup> will be dependent upon the linkage being formed. General methods for preparing the linkage are described, for example, in EP-A 0 372 486 and EP-A 0 381 033 and EP-A 0 478 363, which are incorporated herein by reference.

For instance, if X is CONH, L<sup>1</sup> may be -NH<sub>2</sub> and L<sup>2</sup> may be OH (as in an acid) or Cl (as in an acid chloride). When L<sup>2</sup> is OH, a coupling agent is used.

Similarly, if X is NHCO, L<sup>1</sup> may be -CO<sub>2</sub>H or CO-Cl and L<sup>2</sup> may be -NH<sub>2</sub>. Where X is NHSO<sub>2</sub>, L<sup>1</sup> may be SO<sub>2</sub>Cl and L<sup>2</sup> may be -NH<sub>2</sub>. Where X is SO<sub>2</sub>NH, L<sup>1</sup> may be -NH<sub>2</sub> and L<sup>2</sup> may be SO<sub>2</sub>Cl. Methods to prepare such sulfonyl chlorides are disclosed, for instance, in *J. Org. Chem.*, 23, 1257 (1958).

If X is CH=CH, L<sup>1</sup> may be -CHO and L<sup>2</sup> may be CH=P-Ph<sub>3</sub>. Alternately, L<sup>1</sup> may be CH=P-Ph<sub>3</sub> and L<sup>2</sup> may be CHO.

Where X is CH<sub>2</sub>CH<sub>2</sub> may be obtained by reduction of a suitably protected compound wherein X is CH=CH.

Where X is C≡C, L<sup>1</sup> may be -C≡CH and L<sup>2</sup> may be -Br. Alternately, when X is C≡C, L<sup>1</sup> may be Br, I or CF<sub>3</sub>SO<sub>3</sub> and L<sup>2</sup> may be C≡CH and the coupling may be catalyzed by palladium and a base.

Where X is C(O)CH<sub>2</sub>, L<sup>1</sup> may be C(O)CH<sub>2</sub>Br and L<sup>2</sup> may be hydrogen on a nitrogen of a piperazine ring.

Compounds wherein X is CHOCH<sub>2</sub> may be prepared from a suitably protected compound where X is CH=CH by the procedure disclosed in *J. Org. Chem.*, 54, 1354 (1989).


Compounds wherein X is CH<sub>2</sub>CHOH may be obtained from a suitably protected compound where X is CH=CH by hydroboration and basic oxidation as disclosed in *Tet. Lett.*, 31, 231 (1990).

Scheme I




1


2



3



1



4


5

a) BocO, CH<sub>2</sub>Cl<sub>2</sub>; b) H, Pd/C, EtOH; c) CF<sub>3</sub>CO<sub>2</sub>O, Et<sub>3</sub>N, CH<sub>2</sub>Cl<sub>2</sub>

Scheme I provides a method to prepare certain precursors to be employed in the preparation of compounds described by Formula (I), where Het<sup>1</sup> is a piperidine or piperazine and Het<sup>2</sup> is an aryl group. Accordingly, the terminal secondary nitrogen in a piperazinyl/nitroaryl compound is protected with an appropriate acyl-

type protecting group and the nitro group is reduced to a primary amine to give compounds such as 3 or 4.

Scheme II




9

a)  $\text{CbzCl}$ , 5N  $\text{NaOH}$ ,  $\text{THF}/\text{H}_2\text{O}$ ; b)  $\text{PhSO}_2\text{Cl}$ ,  $\text{tBuOH}$ ,  $\text{Py}$ ; c)  $\text{H}_2$ , Pearlman's catalyst,  $\text{AcOH}$ ; d)  $\text{Boc}_2\text{O}$ ,  $\text{CH}_3\text{CN}$ ; e)  $\text{Bn-Br}$ ,  $\text{Et}_3\text{N}/\text{DMF}$ ; f)  $\text{TFA}$ ,  $\text{CH}_2\text{Cl}_2$ ; g)  $\text{ClCH}_2\text{CO}_2\text{Me}$ ,  $\text{Et}_3\text{N}/\text{CH}_3\text{CN}$ ; h)  $\text{TFA}$ ,  $\text{CH}_2\text{Cl}_2$ ; i)  $\text{Cl-CH}_2\text{CO}_2\text{Et}$ ,  $\text{Et}_3\text{N}/\text{CH}_3\text{CN}$ ; j)  $\text{TFA}$ ,  $\text{CH}_2\text{Cl}_2$ ; k)  $\text{Cl-CH}_2\text{CO}_2\text{tBu}$ ,  $\text{Et}_3\text{N}/\text{CH}_3\text{CN}$ ; l)  $\text{H}_2$  /  $\text{Pd-C}$ ,  $\text{MeOH}$ ; m)  $\text{HCl}$  /  $\text{MeOH}$ ; n)  $\text{ClCH}_2\text{CO}_2\text{tBu}$ ,  $\text{K}_2\text{CO}_3$ ,  $\text{DIEA}$  / acetone; 1 N  $\text{NaOH}$ ,  $\text{H}_2\text{O}$

Scheme II provides a method to prepare certain other precursors to be employed in the preparation of compounds described by Formula (1), where A<sup>1</sup> is CH and A<sup>2</sup> is N, and Q is a single bond. Accordingly, the t-butyl ester of a piperidinyl-4-carboxylic acid, such as compound 2, is prepared from the Cbz-protected derivative, such as compound 6, while the benzyl ester, such as compound 2, is prepared from the Boc-protected derivative, such as compound 8. Such esters are then alkylated with haloacetic acid esters to give, after deprotection of the 4-carboxylic acid ester, 4-carboxy-piperidineacetic esters, such as compounds 10-12.

Scheme III




a)  $\text{SOCl}_2$ ,  $\text{CH}_2\text{Cl}_2$ ; b)  $\text{SOCl}_2$ ,  $\text{CH}_2\text{Cl}_2$

Scheme III provides a method to prepare the acid halides of the precursors in Scheme 2 to be employed in the preparation of compounds described by Formula (1), where A<sup>1</sup> is CH and A<sup>2</sup> is N, and Q is a single bond.


10

Scheme IVa




10. R = Me

12. R = tBu



15a. R = tBu

15b. R = Me




Ex. 1. R = Me

Ex. 2. R = H


- 5 a) R = tBu: HBTU, Et<sub>3</sub>N, DMF; R = Me: DCC / HOBT, Et<sub>3</sub>N, DMF; b) TFA, CH<sub>2</sub>Cl<sub>2</sub>




Scheme IVb



3



16



Ex. 3

- 5 a) EDC / HOBt, Et<sub>3</sub>N, DMF; b) 4M HCl, dioxane, CH<sub>2</sub>Cl<sub>2</sub>

Scheme IVa and IVb also provide a method for the preparation of ester derivatives of compounds described by Formula (1), where Het<sup>1</sup> is a piperazine and Het<sup>2</sup> is an aryl group, X is an amide group, A<sup>1</sup> is CH and A<sup>2</sup> is N, and Q is a single bond. Accordingly, removal of only the amine protecting group from the protected product amide, as exemplified by compound 13b in Scheme IVa and compound 16 in Scheme IVb, affords the title compounds of Example 2 and 3.


13

Scheme V



1


17




18



8



19



Ex. 4

- 5 a) EtO<sub>2</sub>CCl, Et<sub>3</sub>N / CH<sub>2</sub>Cl<sub>2</sub>; b) H<sub>2</sub> 45 psi / 10% Pd-C, EtOH / DMF; c) EDC / HOBt, Et<sub>3</sub>N / DMF; d) 4 M HCl / Dioxane; e) Br-CH<sub>2</sub>CO<sub>2</sub>Me, Et<sub>3</sub>N / DMF


Scheme V provides a method for preparing N-alkoxycarbonyl ester derivatives of compounds described by Formula (1), where Het<sup>1</sup> is a piperazine and Het<sup>2</sup> is an aryl group, X is an amide group, A<sup>1</sup> is CH and A<sup>2</sup> is N, and Q is a

14


single bond. Accordingly, N-alkoxycarbonyl derivatives of piperazinylnitroaryl compounds, such as compound 1, are prepared by treating compounds, such as compound 1, with an alkylchloroformate, the nitro group is reduced by catalytic reduction to a primary amine, and acylated with an N-protected piperidine-4-carboxylic acid, such as compound 8. The N-protecting group is removed by methods known in the art, and alkylated with a haloalkylcarboxylic acid ester, such as methylbromoacetate to give compounds such as the title compound of Example 4.

5

Scheme VI



Ex. 4



Ex. 5

5 a) 1N NaOH, MeOH / THF


Scheme VI provides a method for preparing N-alkoxycarbonyl acid

derivatives of compounds described by Formula (1), where Het<sup>1</sup> is a piperazine and Het<sup>2</sup> is an aryl group, X is an amide group, A<sup>1</sup> is CH and A<sup>2</sup> is N, and Q is a single bond. Accordingly, the esters described in Scheme 5 are saponified by methods known in the art to give compounds such as the title compound of Example 5.

10


5.

Scheme VII



17

Scheme VII, continued



23

Ex. 8

- 5 a) EDC / HOBT, Et<sub>3</sub>N, DMF; b) 4 M HCl / Dioxane; c) Br-CH<sub>2</sub>CO<sub>2</sub>tBu, Et<sub>3</sub>N / DMF; d) 1N NaOH, MeOH / THF; e) 77% aq. CH<sub>2</sub>O, H<sub>2</sub>, 35 psi / 10% Pd-C; f) TFA / CH<sub>2</sub>Cl<sub>2</sub>




Scheme VII provides a method for preparing N-alkyl derivatives of

- 10 compounds described by Formula (1), where Het<sup>1</sup> is a piperazine and Het<sup>2</sup> is an aryl group, X is an amide group, A<sup>1</sup> is CH and A<sup>2</sup> is N, and Q is a single bond. Accordingly, a N-protected piperazinylamine, such as 4 (prepared in Scheme I) is condensed with N-protected piperidine-4-carboxylic acid (prepared in Scheme II) by methods known in the art. The piperidinyl protecting group is removed standard conditions to allow for alkylation with an ester of a haloacetic acid, to give a compound such as 21. Removal of the N-piperazinyl protecting group allows then for alkylation, e.g. reductive alkylation with an aldehyde and an appropriate reducing agent to give, after saponification, N-alkyl-piperazines, such as the title compound of Example 6.

20


18

Scheme VIII




19


Scheme VIII, continued



28



31



Ex. 7



5 a) heat, 48 h; b) 50% NaOH; c)  $\text{SOCl}_2$ ,  $\text{CH}_2\text{Cl}_2$ ; d) DCC / HOBt /  $\text{NH}_2$  /  $\text{NH}_2$  (g), DMF; e) DPPA; f)  $\text{Et}_3\text{N}$ ,  $\text{CH}_2\text{Cl}_2$ ; g) 1N NaOH, MeOH / acetone

10 Scheme VIII provides an alternate method from Scheme VII for preparing N-alkyl derivatives of compounds described by Formula (1), where Het<sup>1</sup> is a piperazine or piperazine and Het<sup>2</sup> is an aryl group, X is an amide group in the reverse sense of compounds prepared in Scheme VII, A<sup>1</sup> is CH and A<sup>2</sup> is N, and Q is a single bond. Accordingly, a mono-alkylpiperazine is treated with a 4-cyanoaryl fluoride to give an alkylpiperazinylcyanide, such as compound 26. Under aqueous alkaline conditions the corresponding arylcarboxylic acid may be obtained. The acid halide


20

of such a compound is condensed with a 4-aminopiperidinylacetic acid ester, such as 30 employing standard acylation conditions to give, after saponification of the ester product, compounds such as the title compound of Example 7. The intermediate compounds such as 30 may be obtained from the corresponding 4-carboxypiperidinylacetic acid esters (prepared in Scheme II) by Curtius rearrangement of the primary amide, such as 29.


Scheme IX



34



35




Ex. 8.

a)  $H_2$  50 psi / 10% Pd-C, DMF; b) EDC / HOBT, DIEA, DMF; c)  $H_2$  47 psi /  $PO_2$ , HCl, MeOH; d) 1N NaOH, EtOH / THF

Scheme IX provides a method for preparing 4-piperidinyl-4-amidopiperidineacetic acids described by Formula (I), where Het<sup>1</sup> is piperazine and Het<sup>2</sup> is an aryl group, X is an amide group, A<sup>1</sup> is CH and A<sup>2</sup> is N, and Q is a single bond. Accordingly, a 4-pyridinylaniline, such as compound 33, is obtained by catalytic reduction of the corresponding nitroaryl compound. The aniline is condensed with a 4-carboxypiperidinylacetic acid ester (prepared in Scheme II) employing a standard set of acylation conditions to give a pyridinylamidide, such as compound 34. Catalytic reduction of such a pyridine affords the corresponding piperidine, which is saponified to give compounds such as the title compound of Example 8.

Scheme X




Ex. 9


a) EDC / HOBT, DIEA, DMF; b) TFA,  $CH_2Cl_2$

Scheme X provides a method for preparing 4-piperidinyl-4-amidopiperidineacetic acids described by Formula (I), where Het<sup>1</sup> is pyridine and Het<sup>2</sup> is an aryl group, X is an amide group, A<sup>1</sup> is CH and A<sup>2</sup> is N, and Q is a single bond. Accordingly, a 4-pyridinylaniline, such as compound 33, is condensed with the relevant 4-carboxypiperidine acetic acid ester (prepared in Scheme II), employing methods known in the art, followed by deesterification to give compounds such as the title compound of Example 9.

Scheme XI



25



26

Scheme XI provides a method for preparing 4-piperazinyl[aryl and heteroaryl]-4-amidopiperidineacetic acids described by Formula (1), where Het<sup>1</sup> is piperazine and Het<sup>2</sup> is an aryl or heteroaryl group. X is an amide group, A<sup>1</sup> is CH and A<sup>2</sup> is N, and Q is a single bond. Accordingly, an N-protected-4-nitroarylpiperazine, such as compound 39, is obtained by arylation of an N-protected piperidine with the appropriate 4-nitro(aryl or heteroaryl) halide, such as compound 38. Catalytic reduction of the nitro compound gives the corresponding aniline, which is condensed with an acylhalide derivative of 4-carboxypiperidine acetic acid ester (prepared in Scheme III), such as compound 8. Protecting group removal by methods known in the art affords the desired compounds, as exemplified by the title compounds of Examples 10 and 11.

10

15

Scheme XII



43


44



45




46




47


11



48



49



Ex. 15

- a)  $\text{BuBr}$ ,  $\text{Et}_3\text{N}$ , DMF; b) 4M  $\text{HCl}$  / dioxane,  $\text{CH}_2\text{Cl}_2$ ; c)  $\text{KNO}_2$ ,  $\text{HCl}$  /  $\text{HOAc}$ ; d)  $\text{LiAlH}_4$ , THF; e) EDC /  $\text{HOBt}$ , DIEA, DMF; f)  $\text{H}_2$ , 50 psi / Pearlman's catalyst, EtOH; g) in  $\text{NaOH}$ , EtOH


Scheme XII provides a method for preparing 4-[N-4'-

(bipiperidinyl)aminocarbonyl-1-piperidineacetic acids described by Formula (1), where  $\text{Het}^1$  is a piperidine and  $\text{Het}^2$  is a piperidine, X is an amide group, A<sup>1</sup> is CH and A<sup>2</sup> is N, and Q is a single bond. Accordingly, a mono-N-benzylbipiperidine, such as compound 45, is prepared by alkylation of a mono-N-protected-bipiperidine, followed by N-protecting group removal. The corresponding N-nitroso compound, such as compound 46, obtained for example by treating the amine with potassium nitrite in a mixture of concentrated hydrochloric and acetic acid, is reduced to the





hydrazine, such as compound 42, with an agent such as  $\text{LiAlH}_4$ . Condensation with a 4-carboxypiperidine acetic acid ester (prepared in Scheme II) by methods known in the art affords, after protecting group removal, monoacylhydrazines such as the title compound in Example 15.


Scheme XIII



Scheme 13, continued



Ex. 16




Ex. 17

a) 1,4-DABCO, DMSO; b) H<sub>2</sub> 40 psi / 10% Pd-C, MeOH; c) DIEA, CH<sub>2</sub>Cl<sub>2</sub>; d) gl. AcOH; e) TFA, CH<sub>2</sub>Cl<sub>2</sub>; f) 1 N NaOH, MeOH / H<sub>2</sub>O semi-prep hplc

Scheme XIII provides a method for preparing 4-piperazinylheteroaryl-4-amidopiperidineacetic acids described by Formula (1), where Het<sup>1</sup> is piperazine and Het<sup>2</sup> is a bicyclic heteroaryl group, X is a covalent bond, A<sup>1</sup> is CH and A<sup>2</sup> is N, and Q is a single bond. Accordingly, a 4-piperazinyl-1,2-diamine bearing a piperazinyl protecting group, such as compound 52, is obtained by arylation of an N-protected piperidine with a 5-chloro-2-nitroaniline, such as 52, and catalytic reduction of the nitro group. Acylation with an acylhalide derivative of a 4-carboxypiperidine acetic acid ester (prepared in Scheme III) gives a mixture of mono- and di-acyl compounds, such as 53-55, which is heated in the presence of glacial acetic acid to give the protected desired bicyclic compound, such as 56. Protecting group removal, followed by semi-preparative chromatography affords the desired bicyclic compound, such as the title compound of Example 16.

Scheme XIII also provides a method for preparing piperazinyl-3,4-bis(amidopiperidineacetic acids) described by Formula (1), where Het<sup>1</sup> is piperazine and Het<sup>2</sup> is an aryl group, X is a covalent bond, m = 2, A<sup>1</sup> is CH and A<sup>2</sup> is N, and Q is a single bond. Accordingly, the semi-preparative chromatography of deprotected product also affords the desired piperazinyl-3,4-bis(amidopiperidineacetic acid), such as the title compound of Example 17.

Scheme XIV



Ex. 18

a) EtOCHO; b)  $\text{BH}_3 \cdot \text{Me}_2\text{S}$ , THF; c)  $\text{Et}_3\text{N}$ ,  $\text{CH}_2\text{Cl}_2$ ; d) IN NaOH, MeOH / THF

5


Scheme XIV provides a method for preparing 4-piperazinylaryl-4-alkylamidopiperidineacetic acids described by Formula (I), where Het<sup>1</sup> is piperazine and Het<sup>2</sup> is an aryl group, X is an alkylamide group, A<sup>1</sup> is CH and A<sup>2</sup> is N, and Q is a single bond. Accordingly, the appropriate N-piperazine-protected, 4-N-alkylaminoaryl/piperazine, such as compound 59, is obtained by reduction of the formyl derivative of an aniline, such as compound 4. Acylation with an acylhalide derivative of a 4-carboxypiperidine acetic acid ester (prepared in Scheme III), followed protecting group removal affords the N-alkylamides, such as the title compound of Example 18.

10


33

34

Scheme XV



Ex. 7




Ex. 19


5 a) H<sub>2</sub> 50 psi/ 10% Pd-C, MeOH

Scheme XV provides a method for preparing 4-piperazinyl-4-amidopiperidineacetic acids described by Formula (I), where Het<sup>1</sup> is piperazine and Het<sup>2</sup> is an aryl group. X is an amide group in the reverse sense of the compounds obtained in Scheme 1. A<sup>1</sup> is CH and A<sup>2</sup> is N, and Q is a single bond. Accordingly, the N-alkyl group in the product obtained from Scheme VIII, where N-alkyl is N-benzyl, is removed by catalytic hydrogenation, to afford compounds such as the title compound of Example 19.

Scheme XVI




Scheme XVI, continued



37

Scheme XVI, continued



Ex. 20

- 5 a)  $\text{SOCl}_2$ , MeOH; b) Piperazine, DMSO; c)  $\text{Boc}_2\text{O}$ ,  $\text{CH}_3\text{CN}$ ; d) IN NaOH, MeOH / THF; e)  $\text{BnBr}$ ,  $\text{Et}_3\text{N}$  / DMF; f) TFA; g) EDC / HOBT /  $\text{NH}_3$  (h) DPPA, DMF; i) DCC / HOBT, DMF /  $\text{Et}_3\text{N}$ ; j)  $\text{H}_2$ , Pearlman's catalyst, MeOH; k)  $\text{Br-CH}_2\text{CO}_2\text{tBu}$ ,  $\text{Et}_3\text{N}$  / DMF; l) Lawesson Reagent, toluene; m)  $\text{NaBH}_4$  /  $\text{NiCl}_2 \cdot 6\text{H}_2\text{O}$ , THF / MeOH; n) 1. Ac-Cl,  $\text{Et}_3\text{N}$  /  $\text{CH}_2\text{Cl}_2$ ; o) TFA /  $\text{CH}_2\text{Cl}_2$

10

- 15 Scheme XVI provides a method for preparing 4-piperazinylaryl-4-methyl(N-acyl)amino-piperidineacetic acids described by Formula (1), where Het<sup>1</sup> is piperazine and Het<sup>2</sup> is an aryl group, X is an N-alkyl-N-acylamino group, A<sup>1</sup> is CH and A<sup>2</sup> is N, and Q is a single bond. Accordingly, a piperazine undergoes arylation upon heating with a 4-fluorobenzoic acid ester. The resulting arylpiperazine is protected with a standard N-protecting group and saponified to the corresponding benzoic acid, such as compound **65**. The benzoic acid is condensed with a piperazinyl-N-protected-4-aminopiperidine, such as compound **68**, to give the corresponding amide. Removal of the piperidine protecting group, and alkylation with a haloacetic acid ester affords a compound such as **70**.


20

The 4-aminopiperidines, such as **68**, are obtained by Curtius rearrangement of the appropriate N-protected piperidine-4-carboxylic acid amide, derived from N-protection of a piperidine-4-carboxylic acid ester, such as compound **6**.

38

The thioamide of a compound, such as 71, that is obtained by treating the amide, such as 74 with the Lawesson reagent, is reduced to the alkylamine, such as 72, via nickel-catalyzed hydroboration. Acylation, followed by protecting group removal affords the desired compound, exemplified by the title compound of Example 20.

Scheme XVII



5


Ex. 21

a) TFA, CH<sub>2</sub>Cl<sub>2</sub>

- 10 Scheme XVII provides a method for preparing 4-piperazinyl-4-thioamido-piperidineacetic acids described by Formula (1), where Het<sup>1</sup> is piperazine and Het<sup>2</sup> is an aryl group, X is an thioamide group, A<sup>1</sup> is CH and A<sup>2</sup> is N, and Q is a single bond. Accordingly, the protecting groups on the thioamides, such as 2, described in Scheme XVI, are removed by methods known in the art to give the desired thioamides, as exemplified by the title compound of Example 21.

15

Scheme XVIII




Ex. 22

a) TFA, CH<sub>2</sub>Cl<sub>2</sub>

Scheme XVIII provides a method for preparing 4-piperazinyl-4-methylamino-piperidineacetic acids described by Formula (1), where Het<sup>1</sup> is piperazine and Het<sup>2</sup> is an aryl group, X is an N-methylamino group, A<sup>1</sup> is CH and A<sup>2</sup> is N, and Q is a single bond. Accordingly, the protecting groups on the preparing 4-piperazinyl-4-methylamino-piperidineacetic acids, such as compound 74, described by Scheme 16 are removed by methods known in the art to give the desired methylamino compounds, as exemplified by the title compound of Example 22.

Scheme XIX




Ex. 23 and 24

5 Ex. 23:

R = CF<sub>3</sub>; X = Na) Cl<sub>3</sub>COCOCi, Et<sub>3</sub>N, dioxane; b) THF; c) 1N NaOH, MeOH

Ex. 24:

10 R = tBuO, X = CH


a) diphosgene, Et<sub>3</sub>N, dioxane; b) Et<sub>3</sub>N, THF; c) 1. TFA, CH<sub>2</sub>Cl<sub>2</sub>, 2. 1N NaOH, EtOH;d) H<sub>2</sub> 50 psi / PtO<sub>2</sub>, HCl, MeOH

15

ZZ X = CH

Scheme XIX provides a method for preparing ureas described by Formula (I), where Het<sup>1</sup> is piperazine and Het<sup>2</sup> is an aryl group, X is an amide group, A<sup>1</sup> is N and A<sup>2</sup> is N or CH, and Q is a single bond. Accordingly, an isocyanate, such as compound 76, prepared from an acyl-piperazinylamine, such as compound 75, is treated with a piperazineacetic acid ester (prepared in Scheme II) to give the protected urea. Protecting group removal affords the desired compound as exemplified in the title compounds of Example 23 and 24. When X = CH, the intermediate, such as compound 77, may be obtained from reduction of a pyridineacetic acid ester, such as 78.

Scheme XX




Ex. 25 and Ex. 28





Scheme XXI also provides a method for preparing 4-[4-(piperazinyl)amido]-cyclohexane-1-acetic acids described by Formula (I), where Het<sup>1</sup> is piperazine and Het<sup>2</sup> is an aryl group, X is an amide group, A<sup>1</sup> is CH and A<sup>2</sup> is CH<sub>3</sub>, and Q is a single bond. Accordingly, the 4-triflate prepared from a 4-hydroxyphenylacetic acid ester is treated with a mixture of palladium(bis)acetate, potassium acetate and 1,1'-bis(diphenylphosphine)ferrocene in an atmosphere of carbon monoxide. Acid quench of the mixture affords the 4-carboxyphenylacetic acid ester, as exemplified by compound 85. Catalytic reduction affords the 4-carboxy-cyclohexanecarboxylic acid ester, which is condensed with a 4-acylpiperazinylaniline (prepared in Scheme I) by methods known to the art to provide the protected amide, such as compound 87. Protecting group removal affords the desired compound as exemplified in the title compound of Example 27.

Scheme XXII



Ex. 28

5 a) 4 M HCl / Dioxane; b) Br-CH<sub>2</sub>CH<sub>2</sub>-CO<sub>2</sub>Me, Et<sub>3</sub>N / DMF; c) IN NaOH, MeOH

Scheme XXII provides a method for preparing 4-[4-(piperazinyl)amido]-piperidine-alkanoic acids described by Formula (I), where Het<sup>1</sup> is piperazine and Het<sup>2</sup> is an aryl group, X is an amide group, A<sup>1</sup> is CH and A<sup>2</sup> is N, and Q is a single bond. Accordingly, the acyl protecting group in the acylpiperidine exemplified by compound 20 (prepared in Scheme VII) is removed by methods known in the art, and alkylated with a haloalkanoic acid ester to provide the piperidinealkanoic acid ester exemplified by compound 88. Protecting group removal affords the desired compound, as exemplified by the title compound in Example 28.

Coupling reagents as used herein denote reagents which may be used to form peptide bonds. Typical coupling methods employ carbodiimides, activated anhydrides and esters and acyl halides. Reagents such as EDC, DCC, DPPA, PPA,

5 BOP reagent, HOBt, N-hydroxysuccinimide and oxalyl chloride are typical. Coupling methods to form amide bonds are generally well known to the art.

The methods of peptide synthesis generally set forth by Bodansky *et al.*, THE PRACTICE OF PEPTIDE SYNTHESIS, Springer-Verlag, Berlin, 1984, Ali *et al.* in *J. Med. Chem.*, 29, 984 (1986) and *J. Med. Chem.*, 30, 2291 (1987) are generally illustrative of the technique and are incorporated herein by reference.

10 Solution synthesis for the formation of amide or peptide bonds is accomplished using conventional methods used to form amide bonds. Typically, the amine or aniline is coupled via its free amino group to an appropriate carboxylic acid substrate using a suitable carbodiimide coupling agent, such as N,N'-dicyclohexyl carbodiimide (DCC), optionally in the presence of catalysts such as 1-hydroxybenzotriazole (HOBt) and dimethylamino pyridine (DMAP). Other methods, such as the formation of activated esters, anhydrides or acid halides, of the free carboxyl of a suitably protected acid substrate, and subsequent reaction with the free amine of a suitably protected amine, optionally in the presence of a base, are also suitable. For example, a protected Boc-amino acid or Cbz-amidino benzoic acid is treated in an anhydrous solvent, such as methylene chloride or tetrahydrofuran (THF), in the presence of a base, such as N-methyl morpholine, DMAP or a trialkylamine, with isobutyl chloroformate to form the "activated anhydride", which is subsequently reacted with the free amine of a second protected amino acid or aniline.

25 The reactive functional groups of the sidechains of each synthetic fragment are suitably protected as known in the art. Suitable protective groups are disclosed in Greene, PROTECTIVE GROUPS IN ORGANIC CHEMISTRY, John Wiley and Sons, New York, 1981. For example, the Boc, Cbz, phthaloyl or Fmoc group may be used for protection of an amino or amidino group. The Boc group is generally preferred for protection of an  $\alpha$ -amino group. A t-Bu, cHex or benzyl ester may be used for the protection of the side chain carboxyl. A benzyl group or suitably substituted benzyl group (e.g., 4-methoxy-benzyl or 2,4-dimethoxy-benzyl) is used to protect

the mercapto group or the hydroxyl group. A suitably substituted carbobenzyloxy group or benzyl group may be also be used for the hydroxyl group or amino group. Suitable substitution of the carbobenzyloxy or benzyl protecting groups is ortho and/or para substitution with chloro, bromo, nitro or methyl, and is used to modify the reactivity of the protective group. Except for the Boc group, the protective groups for the amino moiety are, most conveniently, those which are not removed by mild acid treatment. These protective groups are removed by such methods as catalytic hydrogenation, sodium in liquid ammonia or HF treatment, as known in the art.

10 Acid addition salts of the compounds of this invention are prepared in a standard manner in a suitable solvent from the parent compound and an excess of an acid, such as hydrochloric, hydrobromic, sulfuric, phosphoric, acetic, maleic, succinic or methanesulfonic. The acetate salt form is especially useful. Certain of the compounds form inner salts or zwitterions which may be acceptable. Cationic salts are prepared by treating the parent compound with an excess of an alkaline reagent, such as a hydroxide, carbonate or alkoxide, containing the appropriate cation, or with an appropriate organic amine. Cations such as Li<sup>+</sup>, Na<sup>+</sup>, K<sup>+</sup>, Ca<sup>++</sup>, Mg<sup>++</sup> and NH<sub>4</sub><sup>+</sup> are specific examples of cations present in pharmaceutically acceptable salts.

20 This invention provides a pharmaceutical composition which comprises a compound according to formula (I) and a pharmaceutically acceptable carrier. Accordingly, the compounds of formula (I) may be used in the manufacture of a medicament. Pharmaceutical compositions of the compounds of formula (I) prepared as hereinbefore described may be formulated as solutions or lyophilized powders for parenteral administration. Powders may be reconstituted by addition of a suitable diluent or other pharmaceutically acceptable carrier prior to use. The liquid formulation may be a buffered, isotonic, aqueous solution. Examples of suitable diluents are normal isotonic saline solution, standard 5% dextrose in water or buffered sodium or ammonium acetate solution. Such formulation is especially suitable for parenteral administration, but may also be used for oral administration or contained in a metered dose inhaler or nebulizer for insufflation. It may be desirable to add excipients such as polyvinylpyrrolidone, gelatin, hydroxy cellulose, acacia, polyethylene glycol, mannitol, sodium chloride or sodium citrate.

Alternately, the compounds of this invention may be encapsulated, tableted or prepared in an emulsion or syrup for oral administration. Pharmaceutically acceptable solid or liquid carriers may be added to enhance or stabilize the composition, or to facilitate preparation of the composition. Solid carriers include

- 5 starch, lactose, calcium sulfate dihydrate, terra alba, magnesium stearate or stearic acid, talc, pectin, acacia, agar or gelatin. Liquid carriers include syrup, peanut oil, olive oil, saline and water. The carrier may also include a sustained release material such as glyceryl monostearate or glyceryl distearate, alone or with a wax. The amount of solid carrier varies but, preferably, will be between about 20 mg to about 10 J g per dosage unit. The pharmaceutical preparations are made following the conventional techniques of pharmacy involving milling, mixing, granulating, and compressing, when necessary, for tablet forms; or milling, mixing and filling for hard gelatin capsule forms. When a liquid carrier is used, the preparation will be in the form of a syrup, elixir, emulsion or an aqueous or non-aqueous suspension.
- 15 Such a liquid formulation may be administered directly p.o. or filled into a soft gelatin capsule.

For rectal administration, the compounds of this invention may also be combined with excipients such as cocoa butter, glycerin, gelatin or polyethylene glycols and molded into a suppository.

- 20 The compounds of this invention may be used *in vitro* to inhibit the aggregation of platelets in blood and blood products, e.g., for storage, or for *ex vivo* manipulations such as in diagnostic or research use.

This invention also provides a method of inhibiting platelet aggregation and clot formation in a mammal, especially a human, which comprises the internal administration of a compound of formula (I) and a pharmaceutically acceptable carrier. Indications for such therapy include acute myocardial infarction (AMI), deep vein thrombosis, pulmonary embolism, dissecting aneurysm, transient ischemia attack (TIA), stroke and other infarct-related disorders, and unstable angina.

- 25 Chronic or acute states of hyper-aggregability, such as disseminated intravascular coagulation (DIC), septicemia, surgical or infectious shock, post-operative and post-partum trauma, cardiopulmonary bypass surgery, incompatible blood transfusion, abruptio placenta, thrombotic thrombocytopenic purpura (TTP), snake venom and immune diseases, are likely to be responsive to such treatment. In addition, the
- 30

compounds of this invention may be useful in a method for the prevention of metastatic conditions, the prevention or treatment of fungal or bacterial infection, inducing immunostimulation, treatment of sickle cell disease, and the prevention or treatment of diseases in which bone resorption is a factor.

- 5 The compounds of formula (I) are administered either orally or parenterally to the patient, in a manner such that the concentration of drug in the plasma is sufficient to inhibit platelet aggregation, or other such indication. The pharmaceutical composition containing the compound is administered at a dose between about 0.2 to about 50 mg/kg in a manner consistent with the condition of the patient. For acute therapy, parenteral administration is preferred. For persistent states of hyperaggregability, an intravenous infusion of the peptide in 5% dextrose in water or normal saline is most effective, although an intramuscular bolus injection may be sufficient.
- 10

For chronic, but noncritical, states of platelet aggregability, oral

- 15 administration of a capsule or tablet, or a bolus intramuscular injection is suitable. The compound of this invention is administered one to four times daily at a level of about 0.4 to about 50 mg/kg to achieve a total daily dose of about 0.4 to about 200 mg/kg/day.

This invention further provides a method for inhibiting the reocclusion of an artery or vein following fibrinolytic therapy, which comprises internal administration of a compound of formula (I) and a fibrinolytic agent. It has been found that administration of an peptide in fibrinolytic therapy either prevents reocclusion completely or prolongs the time to reocclusion.

- 20 When used in the context of this invention the term fibrinolytic agent is intended to mean any compound, whether a natural or synthetic product, which directly or indirectly causes the lysis of a fibrin clot. Plasminogen activators are a well known group of fibrinolytic agents. Useful plasminogen activators include, for example, anistreplase, urokinase (UK), pro-urokinase (pUK), streptokinase (SK), tissue plasminogen activator (tPA) and mutants, or variants, thereof, which retain plasminogen activator activity, such as variants which have been chemically modified or in which one or more amino acids have been added, deleted or substituted or in which one or more or functional domains have been added, deleted or altered such as by combining the active site of one plasminogen activator with the
- 25
- 30

fibrin binding domain of another plasminogen activator or fibrin binding molecule. Other illustrative variants include tPA molecules in which one or more glycosylation sites have been altered. Preferred among plasminogen activators are variants of tPA in which the primary amino acid sequence has been altered in the growth factor domain so as to increase the serum half-life of the plasminogen activator. tPA

Growth factor variants are disclosed, e.g., by Robinson *et al.*, EP-A 0 297 589 and Browne *et al.*, EP-A 0 240 334. Other variants include hybrid proteins, such as those disclosed in EP 0 028 489, EP 0 155 387 and EP 0 297 882, all of which are incorporated herein by reference. Anistreplase is a preferred hybrid protein for use in this invention. Fibrinolytic agents may be isolated from natural sources, but are commonly produced by traditional methods of genetic engineering.

Useful formulations of tPA, SK, UK and pUK are disclosed, for example, in EP-A 0 211 592, EP-A 0 092 182 and U.S. Patent 4,568,543, all of which are incorporated herein by reference. Typically the fibrinolytic agent may be formulated in an aqueous, buffered, isotonic solution, such as sodium or ammonium acetate or adipate buffered at pH 3.5 to 5.5. Additional excipients such as polyvinyl pyrrolidone, gelatin, hydroxy cellulose, acacia, polyethylene glycol, mannitol and sodium chloride may also be added. Such a composition can be lyophilized.

The pharmaceutical composition may be formulated with both the compound of formula (I) and fibrinolytic in the same container, but formulation in different containers is preferred. When both agents are provided in solution form they can be contained in an infusion/injection system for simultaneous administration or in a tandem arrangement.

Indications for such therapy include myocardial infarction, deep vein thrombosis, pulmonary embolism, stroke and other infarct-related disorders. The compound of formula (I) is administered just prior to, at the same time as, or just after parenteral administration of tPA or other fibrinolytic agent. It may prove desirable to continue treatment with the peptide for a period of time well after reperfusion has been established to maximally inhibit post-therapy reocclusion. The effective dose of tPA, SK, UK or pUK may be from 0.5 to 5 mg/kg and the effective dose of the compound of this invention may be from about 0.1 to 25 mg/kg.

For convenient administration of the inhibitor and the fibrinolytic agent at the same or different times, a kit is prepared, comprising, in a single container, such

as a box, carton or other container, individual bottles, bags, vials or other containers each having an effective amount of the inhibitor for parenteral administration, as described above, and an effective amount of tPA, or other fibrinolytic agent, for parenteral administration, as described above. Such kit can comprise, for example, both pharmaceutical agents in separate containers or the same container, optionally as lyophilized plugs, and containers of solutions for reconstitution. A variation of this is to include the solution for reconstitution and the lyophilized plug in two chambers of a single container, which can be caused to admix prior to use. With such an arrangement, the fibrinolytic and the compound of this invention may be packaged separately, as in two containers, or lyophilized together as a powder and provided in a single container.

When both agents are provided in solution form, they can be contained in an infusion/injection system for simultaneous administration or in a tandem arrangement. For example, the platelet aggregation inhibitor may be in an i.v. injectable form, or infusion bag linked in series, via tubing, to the fibrinolytic agent in a second infusion bag. Using such a system, a patient can receive an initial bolus-type injection or infusion, of the peptide inhibitor followed by an infusion of the fibrinolytic agent.

The pharmacological activity of the compounds of this invention is assessed by their ability to inhibit the binding of  $^3\text{H}$ -SK&F 107260, a known RGD-fibrinogen antagonist, to the GPIIb/IIIa receptor; their ability to inhibit platelet aggregation, *in vitro*, and their ability to inhibit thrombus formation *in vivo*.

#### Inhibition of RGD-mediated GPIIb-IIIa binding

##### Purification of GPIIb-IIIa

Ten units of outdated, washed human platelets (obtained from Red Cross) were lysed by gentle stirring in 3% octylglucoside, 20 mM Tris-HCl, pH 7.4, 140 mM NaCl, 2 mM  $\text{CaCl}_2$  at 4°C for 2 h. The lysate was centrifuged at 100,000g for 1 h. The supernatant obtained was applied to a 5 mL lentil lectin sepharose 4B column (E.Y. Labs) pre-equilibrated with 20 mM Tris-HCl, pH 7.4, 100 mM NaCl, 2 mM  $\text{CaCl}_2$ , 1% octylglucoside (buffer A). After 2 h incubation, the column was washed with 50 mL cold buffer A. The lectin-retained GPIIb-IIIa was eluted with

buffer A containing 10% dextrose. All procedures were performed at 4°C. The GPIIb-IIIa obtained was >95% pure as shown by SDS polyacrylamide gel electrophoresis.

#### 5 Incorporation of GPIIb-IIIa in Liposomes.

A mixture of phosphatidylserine (70%) and phosphatidylcholine (30%) (Avanti Polar Lipids) were dried to the walls of a glass tube under a stream of nitrogen. Purified GPIIb-IIIa was diluted to a final concentration of 0.5 mg/mL and mixed with the phospholipids in a protein:phospholipid ratio of 1:3 (w:w). The mixture was resuspended and sonicated in a bath sonicator for 5 min. The mixture was then dialyzed overnight using 12,000-14,000 molecular weight cutoff dialysis tubing against a 1000-fold excess of 50 mM Tris-HCl, pH 7.4, 100 mM NaCl, 2 mM CaCl<sub>2</sub> (with 2 changes). The GPIIb-IIIa-containing liposomes were centrifuged at 12,000g for 15 min and resuspended in the dialysis buffer at a final protein concentration of approximately 1 mg/mL. The liposomes were stored at -70°C until needed.

#### Competitive Binding to GPIIb-IIIa

The binding to the fibrinogen receptor (GPIIb-IIIa) was assayed by an indirect competitive binding method using [<sup>3</sup>H]-SK&F-107260 as an ROD-type ligand. The binding assay was performed in a 96-well filtration plate assembly (Millipore Corporation, Bedford, MA) using 0.22 µm hydrophilic durapore membranes. The wells were precoated with 0.2 mL of 10 µg/mL polylysine (Sigma Chemical Co., St. Louis, MO.) at room temperature for 1 h to block nonspecific binding. Various concentrations of unlabeled benzadiazapines were added to the wells in quadruplicate. [<sup>3</sup>H]-SK&F-107260 was applied to each well at a final concentration of 4.5 nM, followed by the addition of 1 µg of the purified platelet GPIIb-IIIa-containing liposomes. The mixtures were incubated for 1 h at room temperature. The GPIIb-IIIa-bound [<sup>3</sup>H]-SK&F-107260 was separated from the unbound by filtration using a Millipore filtration manifold, followed by washing with ice-cold buffer (2 times, each 0.2 mL). Bound radioactivity remaining on the filters was counted in 1.5 mL Ready Solve (Beckman Instruments, Fullerton, CA) in a Beckman Liquid Scintillation Counter (Model LS6800), with 40% efficiency.

Nonspecific binding was determined in the presence of 2 µM unlabeled SK&F-107260 and was consistently less than 0.14% of the total radioactivity added to the samples. All data points are the mean of quadruplicate determinations.

Competition binding data were analyzed by a nonlinear least-squares curve fitting procedure. This method provides the IC<sub>50</sub> of the antagonists (concentration of the antagonist which inhibits specific binding of [<sup>3</sup>H]-SK&F-107260 by 50% at equilibrium). The IC<sub>50</sub> is related to the equilibrium dissociation constant (K<sub>i</sub>) of the antagonist based on the Cheng and Prusoff equation:  $K_i = IC_{50}/(1 + L/K_d)$ , where L is the concentration of [<sup>3</sup>H]-SK&F-107260 used in the competitive binding assay (4.5 nM), and K<sub>d</sub> is the dissociation constant of [<sup>3</sup>H]-SK&F-107260 which is 4.5 nM as determined by Scatchard analysis. 4-[[4-(1-Piperazinyl)phenyl]amino]-carbonyl]-1-piperidineacetic acid, which is a compound of this invention, inhibits [<sup>3</sup>H]-SK&F-107260 binding with a K<sub>i</sub> of about 0.005 µM.

### Inhibition of Platelet Aggregation

Inhibition of platelet aggregation was determined following the procedure described in Nichols, *et al.*, *Thrombosis Research*, 75, 143 (1994). Blood was drawn from the antecubital vein of normal human volunteers who had not taken a cyclooxygenase inhibitor within the previous 14 days into a plastic syringe containing one part 3.8% trisodium citrate to nine parts blood. Platelet rich plasma was prepared by centrifuging the blood at 200 g for 10 min at RT. The platelet rich plasma was drawn off and the remaining blood was centrifuged at 2400 g for 5 min at RT to make platelet poor plasma. Platelet count was measured with a model ZBI Coulter Counter (Coulter Electronics Inc., Hialeah, FL) and was adjusted to 300,000/ $\mu$ l using platelet poor plasma. Platelet aggregation was studied in a Chrono-Log model 400VS Lumi Aggregometer (Chrono-Log, Havertown, PA) using platelet rich plasma stirred at 1200 r.p.m. and maintained at 37°C, with platelet poor plasma as the 100% transmission standard. Concentration-response curves for the ability of compounds to inhibit platelet aggregation, measured as the maximum change in light transmission, induced by a maximal concentration of adenosine diphosphate (10  $\mu$ M) were constructed and the IC<sub>50</sub> was determined as the concentration of antagonist required to produce 50% inhibition of the response to the agonist.

### In Vivo Inhibition of Platelet Aggregation

*In vivo* inhibition of thrombus formation is demonstrated by recording the systemic and hemodynamic effects of infusion of the peptides into anesthetized dogs according to the methods described in Aiken *et al.*, *Prostaglandins*, 19, 629 (1980).

57

The examples which follow are intended to in no way limit the scope of this invention, but are provided to illustrate how to make and use the compounds of this invention. Many other embodiments will be readily apparent and available to those skilled in the art.

5

### Example 1

#### Preparation of 4-[[[4-(1-piperazinyl)phenyl]amino]carbonyl]-1-piperidineacetic acid

10 a) methyl 4-piperidinecarboxylate hydrochloride

Isomipicotic acid (40 g, 310 mmol) was suspended, with stirring, in 250 mL of dry methanol and treated with HCl gas (lecture bottle) until solution was obtained. The solution was then cooled in an ice bath and treatment with HCl gas continued for an additional 20 min. After 2.5 h, the reaction mixture was stripped of solvent using a water aspirator (30°C) and the resulting heavy, white cake suspended in 150 mL dry THF and filtered. The filter cake was washed with two portions of 50 mL THF, followed by washing with 3 x 50 mL Et<sub>2</sub>O, dried briefly in air (suction). The pale-yellow filtrate was discarded. Drying the slightly off-white solid was completed over-night in a desiccator (CaSO<sub>4</sub>) under high vacuum. The title compound (43.25g) was obtained analytically pure. Anal. (C<sub>7</sub>H<sub>14</sub>ClNO<sub>2</sub>) calcd: C, 46.80; H, 7.86; N, 7.80; Cl, 19.73. found C, 46.66; H, 7.78; N, 19.82. TLC R<sub>f</sub> = 0.63 (silica gel, 3: 1 ethanol/conc. aq. ammonium hydroxide).

25 b) 1,1-dimethylethyl-4-(methoxycarbonyl)-1-piperidineacetate

The compound of example 1 (a) (1.8 g, 10 mmol) was vigorously stirred with K<sub>2</sub>CO<sub>3</sub> (1.4 g, 10 mmol) and N,N'-diisopropylethylamine (1.74 mL, 10 mmol) in 60 mL of acetone under a slight positive pressure of argon. To this mixture was added t-butyl chloroacetate (1.6 g, 10 mmol) in 10mL acetone during ~15min and then the reaction mixture was gently refluxed for 5h. After cooling in ice, the mixture was filtered and the solids washed with acetone. The pale yellow filtrate was stripped of solvent and the residue taken up in a mixture of diethylether (100 mL) and water (10 mL). The organic layer was washed successively with water, sat. Na<sub>2</sub>CO<sub>3</sub>, water, and brine, dried (MgSO<sub>4</sub>) and stripped of solvent. Removal of solvent and other volatiles was continued overnight in a constant high vacuum, at 50°C, to yield the title compound as a light yellow liquid (1.2 g) that was analytically pure. Anal.

58

(C<sub>13</sub>H<sub>23</sub>NO<sub>4</sub>) calcd: C, 60.68; H, 9.01; N, 5.44. found: C, 60.28; H, 8.91; N, 5.60. MS (ES), (M+H)<sup>+</sup> = 258.0.

c) sodium N-((4-butoxycarbonyl)methyl)piperidine-4-carboxylate

The compound of example 1 (b) (0.78 g, 3 mmol) was dissolved in 3 mL of methanol and treated with ca. 3 mL 1N NaOH overnight. A small fraction of ester remaining (TLC) was saponified by adding 0.3 mL 1N NaOH and heating in a water bath (50°C) for 2 h, leaving only a trace of the ester (TLC). The reaction mixture was evaporated to dryness on a flash-evaporator, followed by azeotropic drying with toluene and high vacuum at 50°C overnight. The title compound (0.72 g; crude) was used without purification. MS (ES), (M+H)<sup>+</sup> = 244.0. TLC Rf 0.64 (silica gel, 1:1:1:1 ethyl acetate, n-butanol, acetic acid, water).

d) 1-t-butoxycarbonyl-4,4'-nitrophenylpiperazine

1-(4-Nitrophenyl)piperazine (Spectrum Chemical Corp.; 20.7 g, 100 mmol) was suspended in 200 mL of methylene chloride and the dark mixture was stirred in an ice bath under argon. To this cold suspension was added, in portions, solid di-*n*-butyl dicarbonate (26.2 g, 120 mmol). After stirring in ice for a half hour, when vigorous gas evolution subsided and most of the solid had entered solution, the ice bath was removed and stirring continued at rt for an additional 3.5 h. The dark solution was filtered, washed with 4 portions of water (50 mL) and brine (50 mL), filtered through a pad of MgSO<sub>4</sub> and the solvent evaporated *in vacuo*. The title compound is obtained as an orange-yellow solid (27.7 g) showing only trace impurities by TLC. MS (ES), (M+H)<sup>+</sup> = 308.0. Crystallization (26 g) from methanol/water yields analytically pure compound (20 g), mp 144.5-145.5°C. Anal. (C<sub>15</sub>H<sub>21</sub>N<sub>3</sub>O<sub>4</sub>) calcd: C, 58.62; H, 6.89; N, 13.67. found: C, 58.76; H, 6.79; N, 13.59.

e) 1-t-butoxycarbonyl-4,4'-aminophenylpiperazine hydrochloride

The compound of example 1 (d) (0.91 g, 3 mmol) was dissolved in 100 mL 95% ethanol in a Paar vessel, the yellow solution cooled in an ice bath and mixed with 3 mL 1N HCl and 5% palladium on carbon (90 mg). The mixture was hydrogenated on a Paar Shaker (H<sub>2</sub>, 48psi, rt) until after a theoretical amount of hydrogen was consumed in the reaction. The colorless solution was taken to dryness

59

(aspirator) to obtain an off-white solid, which was kept in a high vacuum overnight. The title compound (0.93 g) darkens on exposure to light and air. MS (ES) (M+H)<sup>+</sup> = 278.2; TLC Rf 0.26 (silica gel, 3% methanol/chloroform). f) 4-[[[4-(1-piperazinyl)phenyl]amino]carbonyl]-1-piperidineacetic acid trifluoroacetate

To a stirred, ice-cold suspension of the compound of example 1 (e) (0.5 g, 1.5 mmol) and the compound of example 1 (c) (0.44 g, 1.7 mmol) in 8 mL DMF and 0.42 mL triethylamine (3 mmol) was added solid 2-(1H-benzotriazol-1-yl)-1,3,3-tetramethyluronium hexafluorophosphate and the mixture left stirring in an ice-bath. After 1 h the ice bath was removed, triethylamine (0.42 mL, 3 mmol) was added to bring the mixture to ~ pH 9 and stirring continued overnight at rt, under argon, and for an additional day at 50°C. The reaction mixture was stripped of solvent *in vacuo* and the residue extracted with diethylether, filtered, the ether extract washed with water, sat. NaHCO<sub>3</sub>, and brine, dried over Na<sub>2</sub>SO<sub>4</sub> and the solvent evaporated *in vacuo*. The remaining lt. orange solid (0.5 g) was taken up in 20 mL of methylene chloride and treated with 20 mL TFA, and stirred under argon for 2 h. After the volatiles were removed *in vacuo*, this treatment was repeated and the solid remaining after the solvent was evaporated was dissolved in 10 mL water and purified, portion-wise, by RPHPLC (Vydac C18 support; 0.1% TFA-5% acetonitrile in water). The major component was collected and lyophilized to obtain the title compound as a (hygroscopic) white powder. TLC Rf 0.18 (silica gel; ethyl acetate: n-butanol: acetic acid: water, 1:1:1:1). RPHPLC k' 5.8 (Vydac C18, gradient A: 0.1% TFA-water B 0.08% TFA-acetonitrile, 5-50% acetonitrile during 20 min). MS (ES) (M+H)<sup>+</sup> = 347. H<sup>1</sup> NMR (400 MHz, methanol-d<sub>4</sub>) 7.48 (d, 2H, J = 9 Hz), 7.00 (d, 2H, J = 9 Hz), 4.09 (s, 2H), 3.73 (br s, 2H), 3.35 (s, 8H), 3.15 (br s, 2H), 2.71 (br s, 1H), 2.13 (br s, 4H).

## Example 2

Preparation of Methyl 4-[[[4-(1-piperazinyl)phenyl]amino]carbonyl]-1-piperidine acetate

a) Methyl 4-[[[4-(4-t-butoxycarbonyl-1-piperazinyl)phenyl]amino]carbonyl]-1-piperidine acetate

A mixture of Methyl 4-carboxy-1-acetate (3.0 g, 12.6 mmol), Et<sub>3</sub>N (5.7 mL, 55 mmol), 4-(4-aminophenyl)-1-t-butoxycarbonylpiperazine (3.0 g, 8.6 mmol) in

60



CH<sub>3</sub>CN (50 mL) was treated with HATU (5.0 g, 13.1 mmol). Solution was stirred at RT for 10 hr. Solution was concentrated and residue was precipitated from MeOH/AcOH/H<sub>2</sub>O (20mL/ 0.1 mL/ 60 mL) to give the title compound (3.4g, 75%). MS(ES)<sup>m/e</sup> 461 [M+H]<sup>+</sup>.

5

b) Methyl 4-[[[4-(1-piperazinyl)phenyl]amino]carbonyl]-1-piperidine acetate

A solution of Methyl 4-[[[4-(4-*t*-butoxy carbonyl)-1-

piperazinyl]phenyl]amino]carbonyl]-1-piperidine acetate

(2.1 g, 4.5 mmol) was treated with CH<sub>2</sub>Cl<sub>2</sub>/TFA (10 mL/15 mL) for 20 min.

10 Solution was concentrated and the residue was treated with MeOH/4M HCl (10mL/10 mL). Solution was concentrated until precipitation started. Solution was filtered and the solid was washed with Et<sub>2</sub>O to give the title compound (1.5g, 87%). MS(ES)<sup>m/e</sup> 361 [M+H]<sup>+</sup>. <sup>1</sup>H NMR (250 MHz, DMSO-*d*(6)) δ 1.4-4.5 (m, 22H), 6.9-9.5 (m, 5H). Anal. (C<sub>18</sub>H<sub>23</sub>N<sub>5</sub>O<sub>3</sub> · 2.87HCl (0.5H<sub>2</sub>O) calcd: C, 54.32; H, 7.20; N, 10.29. found: C, 54.52; H, 7.24; N, 10.35.

15

### Example 3

Preparation of Ethyl 4-[[[4-(1-Piperazinyl)phenyl]amino]carbonyl]-1-piperidine acetate 16

20

a) Ethyl 4-[[[4-(1-(*t*-butoxycarbonyl)piperazinyl]phenyl]amino]carbonyl]-1-piperidine acetate 16

EDC (1.69g, 8.8mmole) was added to a solution containing of 4-[1-(*t*-butoxycarbonyl)piperazinyl] aniline 3 (2.22g, 8mmol), HOBt & H<sub>2</sub>O (1.19 g, 8.8 mmole), ethyl 4-carboxypiperidinyl acetate 8 (1.72g, 8mmol) and diisopropylethylamine (1.82mL, 10.4 mmole) in anhydrous DMF (20 mL) at RT. After 20 hr the reaction was concentrated on the rotavap (high vacuum). The residue was dissolved in CH<sub>2</sub>Cl<sub>2</sub> and silica gel chromatography (2 % MeOH/CH<sub>2</sub>Cl<sub>2</sub>) gave the title compound (2.73 g, 72 %): MS (ES)<sup>m/e</sup> 475.2 (M + H)<sup>+</sup>.

25

b) Ethyl 4-[[[4-(1-Piperazinyl)phenyl]amino]carbonyl]-1-piperidine acetate

The compound of 3 (a) (2.6g, 5.48 mmol) was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (30 mL) and treated with HCl (4M in dioxane, 60 mL) for 2 h. The solvent was removed on rotavap and left an off white solid: MS (ES)<sup>m/e</sup> 375.2 (M + H)<sup>+</sup>. Anal. Calcd for C<sub>20</sub>H<sub>26</sub>N<sub>4</sub>O<sub>3</sub> · 43.1 HCl C, 49.27; H, 6.84; N, 11.49; Cl, 22.54. Found: C, 49.16; H, 6.93; N, 11.2; Cl, 22.69.

35

61

### Example 4

Preparation of 4-[[[4-(1-N-ethyloxycarbonyl-4-piperazinyl)phenyl]amino]carbonyl]-1-piperidine-methyl acetate

5

a) 4-[(1-N-ethyloxycarbonyl)piperazinyl]-nitrophenol

Ethylchloroformate (2.5 mL, 24.3 mmol) was added to a cooled solution of 4-piperazinyl nitrophenol (5.0 g, 24.1 mmol) and Et<sub>3</sub>N (6.7 mL, 24.1 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (100 mL). Reaction was stirred at RT for 18 h. Reaction was washed with 5% NaHCO<sub>3</sub> and H<sub>2</sub>O. Organic layer was concentrated to give the title compound (6.27g, 94%) MS(ES)<sup>m/e</sup> 280 [M+H]<sup>+</sup>.

10

b) 4-[(1-N-ethyloxycarbonyl)piperazinyl]-aniline

A mixture of compound of Example 4(a) (3.0 g, 10.8 mmol) and 10% Pd/C (1.0 g) in EtOH/DMF (20 mL/ 30 mL) was hydrogenated at 45 psi for 1 h. Reaction was filtered through Celite and the filtrate was concentrated to afford title compound (2.85 g, 100%). MS(ES)<sup>m/e</sup> 250 [M+H]<sup>+</sup>.

15

c) 4-[[[4-(4-ethyloxycarbonyl)-1-piperazinyl]phenyl]amino]carbonyl]-1-*t*-butoxycarbonyl-piperidine

20

A solution of 4-(4-ethyloxycarbonyl)-1-piperazinylphenylamine (2.7 g, 10.8 mmol), 1-*t*-butoxycarbonyl-4-carboxypiperidine (2.5 g, 10.8 mmol), HOBt (1.6 g, 10.8 mmol) and Et<sub>3</sub>N (2.0 mL, 15 mmol) was treated with EDC (2.1 g, 10.8 mmol). Solution was stirred at RT for 18 hr. Solution was concentrated and the residue was partitioned between EtOAc and H<sub>2</sub>O. EtOAc layer was separated and washed successively with 5% citric acid (2X), H<sub>2</sub>O (1X), 5% NaHCO<sub>3</sub> (2X), H<sub>2</sub>O (1X), brine (1X). EtOAc was dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated to give the title compound (4.69g, 94%). MS(ES)<sup>m/e</sup> 461 [M+H]<sup>+</sup>.

25

d) 4-[[[4-(4-ethyloxycarbonyl)-1-piperazinyl]phenyl]amino]carbonyl]piperidine

30

A solution of compound of Example 4(c) (1.0 g, 2.2 mmol) in 11Cl/Dioxane (4M, 15 mL) was stirred at RT for 1 h. Reaction was concentrated and the residue was treated with Et<sub>2</sub>O. Solution was filtered and the solid was dried *in vacuo* to afford the title compound (0.86g, 100%). MS(ES)<sup>m/e</sup> 361 [M+H]<sup>+</sup>.

35

e) Methyl 4-[[[4-(4-ethyloxycarbonyl)-1-piperazinyl]phenyl]amino]carbonyl]-1-piperidine acetate

62

To a solution of 4-[[[4-(4-ethyloxycarbonyl)-1-

piperazinyl]phenyl]amino]carbonyl]piperidine (0.87 g, 2.2 mmol) and Et<sub>3</sub>N (0.6 mL, 4.4 mmol) in DMF (10 mL) was added methyl bromoacetate (0.208 mL, 2.2 mmol).

After 18h solution was concentrated. Residue was taken up in EtOAc and washed with 5% NaHCO<sub>3</sub>. EtOAc layer was separated and concentrated to give, the title compound (0.40g, 50%). <sup>1</sup>HNMR (250 MHz, DMSO-d<sub>6</sub>) δ 1.0-4.2 (m, 26H), 6.8-7.5 (d, 4H), 9.6 (s, 1H). MS(ES) m/e 433 [M+H]<sup>+</sup>. Anal. (C<sub>24</sub>H<sub>32</sub>N<sub>4</sub>O<sub>3</sub>) calcd: C, 61.09; H, 7.46; N, 12.95. found: C, 61.15; H, 7.53; N, 12.92.

10

### Example 5

Preparation of 4-[[[4-(1-N-ethyloxycarbonyl-4-piperazinyl)phenyl]amino]carbonyl]-1-piperidine acetic acid

15 a) 4-[[[4-(1-N-ethyloxycarbonyl-4-piperazinyl)phenyl]amino]carbonyl]-1-piperidine acetic acid

To a solution of Methyl 4-[[[4-(4-ethyloxycarbonyl)-1-

piperazinyl]phenyl]amino]carbonyl]-1-piperidine acetate

(0.84 g, 1.9 mmol) and MeOH-H<sub>2</sub>O (1/1, 10 mL) was added 1N NaOH (2.2 mL, 2.2 mmol). After 18 hr solution was concentrated. H<sub>2</sub>O was added followed by AcOH to pH 4 (litmus paper). Solution was placed in the refrigerator. Filtration gave the title compound (0.27 g, 33%). <sup>1</sup>HNMR(250MHz,DMSO-d<sub>6</sub>) δ 1.2-4.2 (m, 24H), 6.8-7.6 (dd, 4H), 9.7 (s, 1H). MS(ES) m/e 419 [M+H]<sup>+</sup>. Anal.(C<sub>21</sub>H<sub>29</sub>N<sub>4</sub>·2H<sub>2</sub>O) calcd: C, 55.49; H, 7.54; N, 12.33. found: C, 55.63; H, 7.51; N, 12.29.

25

### Example 6

Preparation of 4-[[[4-(4-Methyl-1-piperazinyl)phenyl]amino]carbonyl]-1-piperidine acetic acid

30

a) t-Butyl-4-[[[4-(4-N-trifluoroacetyl-1-piperazinyl)phenyl]amino]carbonyl]-1-piperidine acetate

A solution of 4-(4-N-trifluoroacetyl-1-piperazinyl)phenyl]amine (0.8 g, 3 mmol), t-butyl-4-carboxy-1-piperidine acetate (0.8 g, 3 mmol), Et<sub>3</sub>N (1.3 mL, 9.0 mmol) in CH<sub>3</sub>CN (25 mL) was treated with HATU (1.25 g, 2.2 mmol). Solution was stirred at RT for 18 hr. Solution was concentrated. Residue was taken up in EtOAc and washed successively with 5% NaHCO<sub>3</sub>, H<sub>2</sub>O. Solution was

63

concentrated and the residue was treated with hot EtOAc. Cooling to RT gave precipitation which was filtered to give the title compound (0.86g, 50%). MS(ES) m/e 499 [M+H]<sup>+</sup>.

5 b) t-Butyl-4-[[[4-(1-piperazinyl)phenyl]amino]carbonyl]-1-piperidine acetate

A solution of t-Butyl-4-[[[4-(4-N-trifluoroacetyl)-1-

piperazinyl]phenyl]amino]carbonyl]-1-piperidine acetate (0.89 g, 1.5 mmol), in

EtOH (5 mL) was treated with 1N NaOH (1.5 mL, 1.5 mmol). After 18 hr solution was concentrated to give the title compound (0.53g, 40%). MS(ES) m/e 289 [M+H]<sup>+</sup>.

10

c) t-Butyl-4-[[[4-(4-methyl-1-piperazinyl)phenyl]amino]carbonyl]-1-piperidine acetate

A stirred mixture of compound of Example 6(b) (0.89 g, 1.49 mmol), Pd/C (5%, 0.1 g) and Formaldehyde (77% soln., 0.15 mL, 1.9 mmol) was hydrogenated at 35 psi for 4 h. Reaction was filtered and the filtrate was concentrated to afford the title compound (0.95g, 80%). MS(ES) m/e 305 [M+H]<sup>+</sup>.

15

d) 4-[[[4-(4-methyl-1-piperazinyl)phenyl]amino]carbonyl]-1-piperidine acetic acid

A solution of t-Butyl-4-[[[4-(4-methyl-1-

piperazinyl)phenyl]amino]carbonyl]-1-piperidine acetate (0.53 g, 1.3 mmol) in 1,4-dioxane (2 mL) was treated with 4M HCl (20 mL). After 3.5 hr solution was concentrated. Residue was purified by prep. HPLC (C-18 reversed phase, 5% CH<sub>3</sub>CN, 0.1% TFA, H<sub>2</sub>O) to give the title compound (0.21g, 50%). MS(ES) m/e 361 [M+H]<sup>+</sup>. <sup>1</sup>HNMR(250MHz,DMSO-d<sub>6</sub>) δ 1.5-4.5 (m, 21H), 6.9-9.5 (m, 5H). Anal.(C<sub>19</sub>H<sub>24</sub>N<sub>4</sub>O<sub>3</sub>·3.25TFA) calcd: C, 42.00; H, 4.01; N, 7.56. found: C, 42.14; H, 4.21; N, 7.86.

25

### Example 7

Preparation of Sodium 4-[[[4-(4-phenylmethyl)-1-piperazinyl]phenyl]carbonyl]amino]-1-piperidine acetate

30

a) 1-benzyl-4-(4-cyanopheno)l)piperazine

A solution of 4-fluorobenzonitrile (3.6 g, 29.7 mmol) and 1-benzylpiperazine (7.5 g, 42.6 mmol) was refluxed for 24 h. Toluene was added and

35

64

washed consecutively with water, brine, dried over  $\text{Na}_2\text{SO}_4$ . Solution was concentrated to give the title compound (0.76g, 10%). MS(ES) m/e 278 [M+H]<sup>+</sup>

b) 1-benzyl-4-(4-carboxyphenyl)piperazine

A solution of compound of Example 7(a) (5 g, 18 mmol) in NaOH solution (51% (w/v) in water) was refluxed for 24 h. Reaction was cooled and treated with HCl (6 N, 80 mL). Reaction was filtered and solid was dried to give title compound (1.9g, 40%). MS(ES) m/e 297 [M+H]<sup>+</sup>.

10 c) 1-benzyl-4-(4-chlorocarbonylphenyl)piperazine

A solution of 1-benzyl-4-(4-carboxyphenyl)piperazine (1.34 g, 3.47 mmol) in  $\text{CH}_2\text{Cl}_2$  (20 mL) was treated with thionyl chloride (1 mL, 35 mmol). Solution was refluxed for 1 hr. Solution was concentrated to an oil which was used as is for the next step (1.9g, Quant.)

15 d) Methyl 4-[[[4-(4-phenylmethyl)-1-piperazinyl]phenyl]carbonyl]amino]-1-piperidine acetate

A cold solution of 1-benzyl-4-(4-chlorocarbonylphenyl)piperazine (1.9 g, 3.5 mmol) and  $\text{Et}_3\text{N}$  (0.5 mL, 3.5 mmol) in  $\text{CH}_2\text{Cl}_2$  (40 mL) was added at once to a cold solution of methyl-4-aminopiperidine acetate (0.7 g, 1.5 mmol) in  $\text{CH}_2\text{Cl}_2$  (20 mL). After 4 hr of stirring at RT solution was filtered. Filtrate was washed successively with 5%  $\text{Na}_2\text{CO}_3$  (2X),  $\text{H}_2\text{O}$  (2X) and dried over  $\text{Na}_2\text{SO}_4$ . Filtration and concentration gave a residue which was purified by flash silica gel chromatography ( $\text{CHCl}_3/\text{MeOH}/\text{Et}_3\text{N}$  95:4:1) to give the title compound (0.47g, 25%). MS(ES) m/e 451 [M+H]<sup>+</sup>.

e) Sodium 4-[[[4-(4-phenylmethyl)-1-piperazinyl]phenyl]carbonyl]amino]-1-piperidine acetate

To a solution of compound of Example 7(d) (0.47 g, 1.04 mmol) in MeOH (5 mL), acetone (5 mL) was added 1N NaOH (1.12 mL, 1.12 mmol). Reaction was heated to 50°C for 40 min. Reaction was cooled and filtered to give the title compound (0.3g, 80%). MS(ES) m/e 437 [M+H]<sup>+</sup>. <sup>1</sup>H NMR (250 MHz, DMSO-d<sub>6</sub>)  $\delta$  1.5-4.5 (m, 21H), 6.8-9.5 (m, 10H). Anal. ( $\text{C}_{23}\text{H}_{31}\text{N}_4\text{O}_3 \cdot 0.5\text{H}_2\text{O}$ ) calcd: C, 63.78; H, 6.80; N, 11.71. found: C, 63.98; H, 6.77; N, 11.92.

#### Example 8

65

#### Preparation of 4-[[[4-(4-piperidinyl)phenyl]amino]carbonyl]-1-piperidine acetate

a) 4-(4-pyridinyl)nitrobenzene 33

The title compound was prepared in 34% yield according to R. Forsyth and F. L. Pyman, J. Chem. Soc. Perkin Trans II, 1926, 2912-24. MS(ESI) m/e 201.0 [M+H]<sup>+</sup>. <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>)  $\delta$  9.05(d, 2H), 8.5 (d, 2H), 8.45 (d, 2H), 8.3 (d, 2H). HPLC k' 5.58 (Ultrasphere  $\Phi$  ODS, gradient, A:acetonitrile B:water-0.1% trifluoroacetic acid, 5-60% acetonitrile during 20 min; UV detection at 220 nm). Anal. ( $\text{C}_{11}\text{H}_8\text{N}_2\text{O}_2 \cdot 3\text{H}_2\text{O}$ ) calcd: C, 45.49; H, 5.2; N, 9.64. Found: C, 45.56; H, 4.99; N, 9.47.

b) 4-(4-pyridinyl)aniline hydrochloride 34

To a solution of Example 8(a) (2.4 g, 10 mmol) in DMF (70 mL) was added 10% Pd/C (0.75 g) under argon. The solution was hydrogenated under 50 psi for 1 h. The solution was filtered through celite, and concentrated to afford the title compound as yellow solid (1.63 g, 79%). MS(ESI) m/e 171.0 [M+H]<sup>+</sup>. TLC R<sub>f</sub> 0.63 (silica gel, 1:1 hexane:ethyl acetate). HPLC k' 1.88 (Ultrasphere  $\Phi$  ODS, gradient, A:acetonitrile B:water-0.1% trifluoroacetic acid, 5-60% acetonitrile during 20 min; UV detection at 220 nm).

c) Ethyl-4-[[[4-(4-pyridinyl)phenyl]amino]carbonyl]-1-piperidine acetate 35

EDC (636 mg, 3.3 mmol) was added to a solution of the compound of Example 8 (b) (682 mg, 3.3 mmol), ethyl-4-carboxypiperidine acetate 14 (646 mg, 3 mmol), HOBT,  $\text{H}_2\text{O}$  (446 mg, 3.3 mmol) and DIEA (1.2 mL, 6.8 mmol) in anhydrous DMF (10 mL) at RT. After stirring for 20 h, the reaction was concentrated on rotavap (high vacuum). The resulting residue was taken into EtOAc and washed successively with  $\text{H}_2\text{O}$  (3 x 20 mL), 10%  $\text{NaHCO}_3$  (2 x 20 mL) and saturated NaCl. The organic extract was dried (anhydrous  $\text{Na}_2\text{SO}_4$ ), filtered and concentrated to yield the title compound (350 mg, 32%) as a yellow solid. HPLC k' 5.3 (Ultrasphere  $\Phi$  ODS, gradient, A:acetonitrile B:water-0.1% trifluoroacetic acid, 5-60% acetonitrile during 20 min; UV detection at 220 nm). MS (ESI) m/e 368.2[M+H]<sup>+</sup>.

d) Ethyl-4-[[[4-(4-piperidinyl)phenyl]amino]carbonyl]-1-piperidine acetate 36

To a solution of the compound of Example 8(c) (260 mg, 0.7 mmol), in methanol (15 mL) was added  $\text{PtO}_2$  (150 mg, 0.66 mmol) and HCl (2 mL of 1N) under argon. The solution was hydrogenated under 47 psi for 4 h. The solution was

66

filtered through celite, concentrated and triturated with ethyl acetate to afford the title compound as off white solid (30 mg). MS(ESI) m/e 374.2 [M+H]<sup>+</sup>. <sup>1</sup>H NMR (400 MHz, DMSO-D<sub>6</sub>) δ 10.25 (brs, 1H), 8.8 (brs, 1H), 7.55 (d, 1H), 7.15 (d, 1H), 4.25 (q, 2H), 3.55 (brs, 2H), 3.3 (brs, 8H), 2.6-3 (m, 4H), 2.0 (m, 4H), 1.9 (m, 4H), 1.25 (t, 3H). HPLC k' 5.0 (Ultrasphere ® ODS, gradient, A:acetonitrile B:water-0.1% trifluoroacetic acid, 5-60% acetonitrile during 20 min; UV detection at 220 nm). Anal. (C<sub>21</sub>H<sub>23</sub>N<sub>3</sub>O<sub>3</sub> · 2 HCl · 0.5H<sub>2</sub>O) calcd: C, 55.38; H, 7.53; N, 9.23. Found: C, 55.16; H, 7.49; N, 9.16.

10 c) 4-[[[4-(4-piperidinyl)phenyl]amino]carbonyl]-1-piperidine acetic acid sodium salt

To a solution of the compound of Example 8(d) (370 mg, 0.83 mmol) in 1:1 ethanol-THF mixture (20 mL) was added a solution of 1N NaOH (2.5 mL, 2.5 mmol) and was stirred at RT for 20 h. The resulting white precipitate was collected by filtration to yield the title compound (170 mg) as white solid. HPLC k' 2.38 (Ultrasphere ® ODS, gradient, A:acetonitrile B:water-0.1% trifluoroacetic acid, 5-60% acetonitrile during 20 min; UV detection at 220 nm). <sup>1</sup>H NMR (400 MHz, DMSO-D<sub>6</sub>) δ 10.05 (s, 1H), 9.9 (brs, 1H), 8.5 (dd, 1H), 7.50 (d, 2H), 7.15 (d, 2H), 4.2 (s, 2H), 3.65 (brd, 2H), 3.3 (br d, 2H), 2.95 (m, 4H), 2.75 (m, 1H), 2.55 (m, 1H), 1.7-2.0 (m, 8H). MS (ESI) m/e 346.0 [M+H]<sup>+</sup>. Anal. (C<sub>19</sub>H<sub>26</sub>N<sub>3</sub>O<sub>4</sub> Na · 0.25 EtOH, 2.25 H<sub>2</sub>O) calcd: C, 55.84; H, 7.69; N, 10.02. Found: C, 55.80; H, 7.56; N, 9.68.

#### Example 9

##### Preparation of 4-[[[4-(4-pyridinyl)phenyl]amino]carbonyl]-1-piperidine acetic acid

a) 1-Butoxycarbonyl-4-[[[4-(4-pyridinyl)phenyl]amino]carbonyl]-1-piperidine acetate 37

EDC (316.3 mg, 1.65 mmol) was added to a solution 4-(4-pyridinyl)aniline hydrochloride 34 (342 mg, 1.65 mmol), t-butoxycarbonyl-4-carboxypiperidine acetate 12 (365 mg, 1.5 mmol), HOBT·H<sub>2</sub>O (223 mg, 1.65 mmol) and DIEA (634 µL, 3.64 mmol) in anhydrous DMF (10 mL) at RT. After stirring for 20 h, the reaction was concentrated on rotavap (high vacuum). The resulting residue was taken into EtOAc and washed successively with H<sub>2</sub>O (3 x 20 mL), 10% NaHCO<sub>3</sub> (2 x 20 mL) and saturated NaCl. The organic extract was dried (anhydrous Na<sub>2</sub>SO<sub>4</sub>), filtered and concentrated to yield the title compound (500 mg, 84%) as a yellow

solid. HPLC k' 6.35 (Ultrasphere ® ODS, gradient, A:acetonitrile B:water-0.1% trifluoroacetic acid, 5-60% acetonitrile during 20 min; UV detection at 220 nm). MS (ESI) m/e 396.2 [M+H]<sup>+</sup>.

5 b) Preparation of 4-[[[4-(4-pyridinyl)phenyl]amino]carbonyl]-1-piperidine acetic acid

The compound of Example 9(a) (0.5 g, 1.29 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (10 mL) was added trifluoroacetic acid (10 mL) at RT and stirred for 5h. The resulting solution was then concentrated to dryness on rotavap and evaporated with CH<sub>2</sub>Cl<sub>2</sub> twice. The resulting oily residue was dissolved in H<sub>2</sub>O, and the pH was adjusted to 8.45 using dilute ammonium hydroxide. The aqueous solution was purified on flash ODS column (step gradient, 2-16% acetonitrile/water. The fractions containing the pure compound were collected, concentrated and lyophilized to yield the title compound (150 mg, 35%) as a white powder. HPLC k' 4.23 (Ultrasphere ® ODS, gradient, A:acetonitrile B:water-0.1% trifluoroacetic acid, 5-60% acetonitrile during 20 min; UV detection at 220 nm). MS (ESI) m/e 340.2 [M+H]<sup>+</sup>. Anal. (C<sub>19</sub>H<sub>21</sub>N<sub>3</sub>O<sub>3</sub> · 2.25 H<sub>2</sub>O) calcd: C, 60.07; H, 6.77; N, 11.06. Found: C, 59.76; H, 6.57; N, 10.88. <sup>1</sup>H NMR (400 MHz, DMSO-D<sub>6</sub>-TFA) δ 10.42 (s, 1H), 9.85 (br s, 1H), 8.88 (dd, 2H), 8.38 (d, 2H), 8.05 (d, 2H), 7.82 (d, 2H), 4.15 (s, 2H), 3.6 (brd, 2H), 3.35-3.05 (brd, 3H), 2.0 (m, 4H).

#### Example 10

##### Preparation of 4-[[[5-(2-(1-piperazinyl)pyridinyl)amino]carbonyl]-1-piperidine acetic acid

a) 2-(1-tert-Butoxycarbonyl)piperazinyl)-5-nitropyridine 40

A mixture of tert-butyl 1-piperazinecarboxylate (1.0 g, 5.3 mmol), 2-chloro-5-nitropyridine (0.85 g, 5.3 mmol) and triethylamine (1.58 mL, 11.0 mmol) in CH<sub>3</sub>CN (50 mL) were stirred at rt temperature for 18 hr. The yellow suspension was concentrated, the residue was taken up in CH<sub>2</sub>Cl<sub>2</sub> (125 mL) and washed with 5% NaHCO<sub>3</sub> (2 x 25 mL) and brine (2 x 25 mL), dried over MgSO<sub>4</sub> and concentrated to dryness in vacuo to yield the yellow titled compound (1.65 g, 100%). MS (ES) m/e 309.2 [m+H]<sup>+</sup>.

35 b) 2-(1-tert-Butoxycarbonyl)piperazinyl)-5-aminopyridine 41

The compound of Example 10(a) (1.65 g, 5.3 mmol) was suspended in MeOH (200 mL) in the presence of 10% Pd/C (0.17 g) and hydrogenated in a Parr

shaker under a hydrogen atmosphere (40 psi) at room temperature for 1.5 h. The catalyst was filtered and the filtrate was concentrated to yield the light-pink titled compound (1.4 g, 95%).

- 5 c) Ethyl 4-[[[5-[2-[[1-(tert-butoxycarbonyl)piperazinyl]pyridinyl]amino]carbonyl]-1-piperidineacetate 42

The compound of Example 10 (b) (1.4 g, 5.0 mmol) and pyridine (1.1 mL) in  $\text{CH}_2\text{Cl}_2$  (20 mL) was added to a suspension of the acid chloride in  $\text{CH}_2\text{Cl}_2$  (10 mL) obtained from refluxing 1-ethoxycarbonylmethyl-4-piperidinecarboxylic acid hydrogen chloride (1.1 g, 4.3 mmol) in  $\text{SOCl}_2$  (10 mL) for 15 min, followed by removal of  $\text{SOCl}_2$ . After 18 h, the reaction mixture was poured into 5%  $\text{NaHCO}_3$  (75 mL) and extracted with  $\text{CH}_2\text{Cl}_2$  (3 x 20 mL). The combined organic fractions were backwashed with 5%  $\text{NaHCO}_3$  (2 x 50 mL) and brine (2 x 50 mL), dried over  $\text{MgSO}_4$  and concentrated. The semi-solid was triturated with EtOAc to give the title compound (1.25 g, 60%). MS (ES) m/e 476.2 [m+H]<sup>+</sup>. Anal. Calcd. for  $\text{C}_{24}\text{H}_{37}\text{N}_5\text{O}_5$ : C, 60.61; H, 7.84; N, 14.73; Found C, 60.58; H, 7.76; N, 14.65.

- 15 d) 4-[[[5-[2-(1-Piperazinyl)pyridinyl]amino]carbonyl]-1-piperidineacetic acid

The compound of Example 10 (c) (650 mg, 1.36 mmol) was dissolved in  $\text{CH}_2\text{Cl}_2$  (45 mL) and treated with 4.0 M HCl in dioxane (6.8 mL) for 2 h. The intermediate HCl salt (510 mg, 100%, MS (ES) m/e 376.2 [m+H]<sup>+</sup>) was obtained by filtration of the resulting suspension, which was dissolved in 0.95 N NaOH solution (8.2 mL, 7.8 mmol) in EtOH (20 mL) and stirred for 18 h. The mixture was concentrated and filtered to give sodium salt of the titled compound (25 mg). MS (ES) m/e 348.2 [m+H]<sup>+</sup>. Anal. Calcd. for  $\text{C}_{17}\text{H}_{24}\text{N}_5\text{O}_3\text{Na}$ : C, 46.83; H, 7.26; N, 16.06; Found C, 47.02; H, 6.56; N, 15.61.

#### Example 11

- 30 Preparation of 4-[[[4-(1-Piperazinyl)-3-chlorophenyl]amino]carbonyl]-1-piperidineacetic acid bis(trifluoroacetate)

- a) 3-Chloro-4-(4-t-butoxycarbonyl-1-piperazinyl)nitrobenzene 44a

A solution of t-butyl 1-piperazinecarboxylate 38 (1.0 g, 5.37 mmole), 3-chloro-4-fluoronitrobenzene (0.94 g, 5.37 mmole) and diisopropylethylamine (0.70 g, 5.4 mmole) in DMSO (30 mL) was heated 16 hours at 100°, cooled to r.t. and poured into ice water (300 mL). The suspension was extracted with ethyl acetate (2

x 60 mL.). The combined organic layers were washed with water (3 x 40 mL.), dried over  $\text{Na}_2\text{SO}_4$  and evaporated to dryness to give an orange solid which was recrystallized twice from methanol to give the title compound, 1.2 g. MS (M+H)<sup>+</sup> 342.0

- b) 3-Chloro-4-(4-t-butoxycarbonyl-1-piperazinyl)aniline 45a

A suspension of the compound of Example 11 (a) (0.34 g, 1.0 mmole) in ethanol (50 mL.) containing 10% Pd on C (0.2 g.) was hydrogenated at 40 psi at r.t. After 40 minutes, the mixture was filtered and evaporated to dryness to give the title compound, 0.31 g. MS (M+H)<sup>+</sup> 312.2

- c) Ethyl 4-[[[4-(4-t-butoxycarbonyl)-1-piperazinyl]-3-chlorophenyl]amino]carbonyl]-1-piperidineacetate 46a

A solution of ethyl 4-(chlorocarbonyl)-1-piperidineacetate hydrochloride 14 (0.27 g, 1.0 mmole) in methylene chloride (10 mL.) was added dropwise to a solution of the compound of Example 11 (b) (0.31 g, 1.0 mmole) and pyridine (0.16 g, 2.0 mmole) in methylene chloride (30 mL.). The solution was stirred 16 hours, extracted with 5%  $\text{Na}_2\text{CO}_3$  (2 x 10 mL.) then with water (10 mL.), dried over  $\text{Na}_2\text{SO}_4$  and evaporated to dryness. The residue was chromatographed (2.5%  $\text{CH}_3\text{OH}/\text{CH}_2\text{Cl}_2$ , silica) to give the title compound, 0.27 g. MS (M+H)<sup>+</sup> 509.4

- d) 4-[[[4-(1-Piperazinyl)-3-chlorophenyl]amino]carbonyl]-1-piperidineacetic acid bis(trifluoroacetate)

A solution of the compound of Example 11 (c) in a mixture of methylene chloride (35 mL.) and trifluoroacetic acid (10 mL.) was kept 16 hours at r.t. Solvents were removed and the residue redissolved in a mixture of methanol (35 mL.), water (3.5 mL.) and 1 N NaOH (3.75 mL.). The solution was heated at 50° for 2 hours, cooled to r.t., quenched with trifluoroacetic acid (1.0 mL.) and evaporated to dryness to give a pale yellow solid which was purified by semi-preparative HPLC (YMC ODS-AQ, 10  $\mu\text{m}$ , 120  $\times$ , 50 X 250 mm.; 13%  $\text{CH}_3\text{CN}/\text{H}_2\text{O}$ , 0.1% TFA; 90 mL./min.) to give the title compound, 0.213 g. MS (M+H)<sup>+</sup> 381.2

#### Example 12

- 35 Preparation of 4-[[[4-(1-piperazinyl)amino]-1-yl]amino]carbonyl]-1-piperidineacetic acid bis(trifluoroacetate)

## a) 4-(4-(4-t-butoxycarbonyl-1-piperazinyl)-1-nitronaphthylene 44b

Using the procedure of Example 11 (a), except substituting 1-fluoro-4-nitronaphthalene 43b for 3-chloro-4-fluoronitrobenzene 43a, the title compound was prepared, 1.2 g. MS (M+H)<sup>+</sup> 358.2

5

## b) 1-Amino-4-(4-(4-t-butoxycarbonyl-1-piperazinyl)-naphthalene 45b

Using the procedure of Example 11 (b), except substituting the compound of Example 12 (a) for the compound of Example 11 (a), the title compound was prepared, 0.65 g. MS (M+H)<sup>+</sup> 338.2

10

## c) Ethyl 4-[[[4-(4-t-butoxycarbonyl-1-piperazinyl)naphth-1-yl]amino]carbonyl]-1-piperidineacetate 46b

Using the procedure of Example 11 (c), except substituting the compound of Example 12 (b) for the compound of Example 11 (b), the title compound was prepared, 0.45 g. MS (M+H)<sup>+</sup> 525.2

15

## d) 4-[[[4-(1-piperazinyl)naphth-1-yl]amino]carbonyl]-1-piperidineacetic acid bis(trifluoroacetate)

Using the method of Example 11 (d), except substituting the compound of Example 12 (c) for the compound of Example 11 (c), the title compound was prepared, 0.407 g. MS (M+H)<sup>+</sup> 397.2

20

## Example 13

## 25 preparation of 4-[[[4-(1-Piperazinyl)-3-(trifluoromethyl)phenyl]amino]carbonyl]-1-piperidineacetic acid bis(trifluoroacetate)

## a) 3-Trifluoromethyl-4-(4-t-butoxycarbonyl-1-piperazinyl)nitrobenzene 44c

Using the procedure of Example 11 (a), except substituting 4-fluoro-3-(trifluoromethyl)nitrobenzene 43c for 3-chloro-4-fluoronitrobenzene 43a, the title compound was prepared, 0.93 g. MS (M+H)<sup>+</sup> 376.2

30

## b) 3-Trifluoromethyl-4-(4-t-butoxycarbonyl-1-piperazinyl)aniline 45c

Using the procedure of Example 11 (b), except substituting the compound of Example 13 (a) for the compound of Example 11 (a), the title compound was prepared, 0.69 g. MS (M+H)<sup>+</sup> 346.2

35

71

## c) Ethyl 4-[[[4-(4-t-butoxycarbonyl-1-piperazinyl)-3-(trifluoromethyl)-phenyl]amino]carbonyl]-1-piperidineacetate 46c

Using the procedure of Example 11 (c), except substituting the compound of Example 13 (b) for the compound of Example 11 (b), the title compound was prepared, 0.45 g. MS (M+H)<sup>+</sup> 543.2

5

## d) 4-[[[4-(1-Piperazinyl)-3-(trifluoromethyl)phenyl]amino]carbonyl]-1-piperidineacetic acid bis(trifluoroacetate)

Using the method of Example 11 (d), except substituting the compound of Example 13 (c) for the compound of Example 11 (c), the title compound was prepared, 0.302 g. MS (M+H)<sup>+</sup> 415.2

10

## Example 14

## 15 preparation of 4-[[[4-(1-Piperazinyl)-2-(trifluoromethyl)phenyl]amino]carbonyl]-1-piperidineacetic acid bis(trifluoroacetate)

## a) 2-Trifluoromethyl-4-(4-t-butoxycarbonyl-1-piperazinyl)nitrobenzene 44d

Using the procedure of Example 11 (a), except substituting 4-fluoro-2-(trifluoromethyl)nitrobenzene 43d for 3-chloro-4-fluoronitrobenzene 43a, the title compound was prepared, 1.58 g. MS (M+H)<sup>+</sup> 376.2

20

## b) 2-Trifluoromethyl-4-(4-t-butoxycarbonyl-1-piperazinyl)aniline 45d

Using the procedure of Example 11 (b), except substituting the compound of Example 14 (a) for the compound of Example 11 (a), the title compound was prepared, 0.69 g. MS (M+H)<sup>+</sup> 346.2

25

## c) Ethyl 4-[[[4-(4-t-butoxycarbonyl-1-piperazinyl)-3-(trifluoromethyl)-phenyl]amino]carbonyl]-1-piperidineacetate 46d

Using the procedure of Example 11 (c), except substituting the compound of Example 14 (b) for the compound of Example 11 (b), the title compound was prepared, 0.34 g. MS (M+H)<sup>+</sup> 543.2

30

## d) 4-[[[4-(1-Piperazinyl)-3-(trifluoromethyl)phenyl]amino]carbonyl]-1-piperidineacetic acid bis(trifluoroacetate)

35

72

Using the method of Example 11 (d), except substituting the compound of Example 14 (c) for the compound of Example 11 (c), the title compound was prepared, 0.172 g. MS (M+H)<sup>+</sup> 415.2

### Example 15

Preparation of 4-[N-(4,4'-Bipiperidinyl) amino] carbonyl-L-piperidine acetic acid

a) N-Boc-N'-benzylbipiperidine 48

Benzylbromide (2.19ml, 18.5 mmol) was added to a mixture containing N-tert-butoxycarbonyl bipiperidine 47 (3.3g, 12.3mmol) and triethylamine (2.57ml, 18.5mmol) in anhydrous DMF (10 ml), and the resulting mixture was stirred at RT for 4hr. The reaction was concentrated on the rotavap (high vacuum). And the residue was dissolved in EtOAc, washed successively with H<sub>2</sub>O, brine, dried (MgSO<sub>4</sub>), concentrated and silica chromatography gave the title compound (2.8g, 64%) as a white solid: MS (ES) m/e 359.2 (M + H)<sup>+</sup>.

b) N-benzyl bipiperidine 47

Following the procedure of Example 3 (b), except substituting the compound of 15 (a) for Ethyl-4-[[[4-(1-(tert-butoxycarbonyl)piperazinyl] phenyl] amino] carbonyl]-1-piperidine acetate 16, the title compound (61%) as a white solid: (ES) m/e 259.2 (M + H)<sup>+</sup>.

c) N-benzyl-N'-nitrosobipiperidine 50

HCl (480μl, 5.75mmol, conc.) was added dropwise to a mixture containing the compound of 15 (b) (1.34g, 4.6mmol) and acetic acid (4.6ml, glacial) with cooling. Then it was heated to 25°C and added a solution of potassium nitrite (1.17g, 13.8mmol) in H<sub>2</sub>O (3ml) and kept at 25°C for another 30min. The product was extracted by CH<sub>2</sub>Cl<sub>2</sub> and washed by Na<sub>2</sub>CO<sub>3</sub>, brine, after drying (MgSO<sub>4</sub>) and concentrating gave 400mg (30% yield) as a white solid: MS (ES) m/e 288.5 (M + H)<sup>+</sup>.

d) N-benzyl-N'-aminobipiperidine 51

To a refluxing LAH (3.5mmol) in THF (7ml) was added dropwise a solution of the compound of Example 15 (c) (400mg, 1.4mmol) in THF (10ml) with stirring over 15 min. The resulting mixture was continued refluxing for 4hr. CH<sub>3</sub>OH (a few drops) and NaOH (1M, 5ml) was added, the product was extracted by THF

and Et<sub>2</sub>O, after drying (MgSO<sub>4</sub>) and concentrating left 310mg (81% yield) as a white solid: MS (ES) m/e 274.2 (M + H)<sup>+</sup>.

e) Ethyl 4-[N-benzylbipiperidinyl] amino] carbonyl-L-piperidine acetate 52

Following the procedure of Example 3 (a), except substituting the compound of 15(d) for 4-[1-(tert-butoxycarbonyl)piperazinyl] aniline 2 the title compound (600mg, 50%) as a white solid: (ES) m/e 471.4 (M + H)<sup>+</sup>.

f) Ethyl 4-[N-(4,4'-Bipiperidinyl) amino] carbonyl-L-piperidine acetate 53

A mixture of the compound of 15 (e) (530mg, 1.13mmol) and Pearlman's catalyst (500mg) in EtOH (100ml) was subjected to hydrogenation at 50psi in a Parr apparatus at RT for 5 hr. The catalyst was removed by filtration and the filtrate was concentrated left 390mg (91%) as a colorless foam: MS (ES) m/e 381.2 (M + H)<sup>+</sup>.

g) 4-[N-(4,4'-Bipiperidinyl) amino] carbonyl-L-piperidine acetic acid:

Following the procedure of Example 8 (c), the compound of Example 15 (d) was saponified to give the title compound as a white powder: MS (ES) m/e 353.5 (M + H)<sup>+</sup>. Anal. Calcd for C<sub>18</sub>H<sub>25</sub>N<sub>4</sub>O<sub>3</sub> 41.5 TFA: H<sub>2</sub>O: C, 46.58; H, 6.61; N, 10.35 Found: C, 46.59; H, 6.84; N, 10.36.

### Example 16

Preparation of 4-[4-(1-Piperazinyl)benzimidazol-2-yl]-1-piperidineacetic acid, tris(trifluoroacetate)

a) 5-[4-(t-butoxycarbonyl)-1-piperazinyl]-2-nitroaniline 55

A solution of t-butyl 1-piperazine carboxylate 38 (1.0 g., 5.4 mmole), 5-chloro-2-nitroaniline 54 (0.93 g., 5.4 mmole) and 1,4-diazabicyclo[2.2.2]octane (0.61 g., 54 mmole) in DMSO (10 mL) was heated 16 hours under argon, cooled to r.t. and diluted with water (150 mL) and extracted with ethyl acetate (3 X 50 mL). The combined organic layers were washed with water (3 X 30 mL) and dried over Na<sub>2</sub>SO<sub>4</sub>. Solvent was removed and the residue chromatographed (40% EtOAc/Hexane, silica) to give the title compound as a yellow solid, (0.5 g., 30%). MS (M+H)<sup>+</sup> 323.

b) 4-[4-(t-butoxycarbonyl)-1-piperazinyl]-o-phenylenediamine 56

A solution of the compound of example 16 (a) (0.30 g., 0.93 mmole) in ethanol (50 mL) containing 10% Pt on carbon (0.10 g.) was hydrogenated at 40 psi

for 40 minutes at r.t. The suspension was filtered and evaporated to give the title compound as a colorless solid, 0.27 g. MS (M+H)<sup>+</sup> 293.2.

- 5 c) Mixture of ethyl 4-[[[4-(4-(t-butoxycarbonyl)-1-piperazinyl)-2-aminophenyl]-amino]carbonyl]-piperidin-4-yl-1-acetate, ) ethyl 4-[[[5-(4-(t-butoxycarbonyl)-1-piperazinyl)-2-aminophenyl]amino]carbonyl]-piperidin-4-yl-1-acetate and N,N'-[4-(1-piperazinyl)-1,2-phenylene]bis-[1-(carboethoxymethyl)-4-piperidinecarboxamide] 57-59

10 A solution of ethyl 4-(chlorocarbonyl)-1-piperidineacetate hydrochloride 14 (0.25 g, 0.93 mmole) in methylene chloride (5.0 mL) was added dropwise over 10 minutes to a solution of the Compound of Example 16b (0.27 g, 0.93 mmole) and diisopropylethylamine (0.26 g, 2.0 mmole) in methylene chloride (10 mL). The mixture was stirred 16 hours, diluted with methylene chloride (35 mL) and extracted with 5% Na<sub>2</sub>CO<sub>3</sub>. The organic layer was washed with water (3 X 20 mL), dried over Na<sub>2</sub>SO<sub>4</sub> and evaporated to give a pale yellow solid which was triturated with ether (30 mL) to give the title mixture (0.17 g). MS (M+H)<sup>+</sup> 489.4, 674.5.

- 20 d) Mixture of ethyl 4-[5-(4-(t-butoxycarbonyl)-1-piperazinyl)benzimidazol-2-yl]-1-piperidineacetate and N,N'-[4-(1-piperazinyl)-1,2-phenylene]bis-[1-(carboethoxymethyl)-4-piperidinecarboxamide] 60-61

A solution of the mixture of Example 16 (c) in glacial acetic acid (30 mL.)

was

25 heated 16 hours at 65°, cooled to r.t. and evaporated to dryness to give the title mixture.

MS (M+H)<sup>+</sup> 471.4, 674.5.

- 30 e) 4-[5-(1-Piperazinyl)benzimidazol-2-yl]-1-piperidineacetic acid, tris(trifluoroacetate) and N,N'-[4-(1-piperazinyl)-1,2-phenylene]bis-[1-(carboxymethyl)-4-piperidinecarboxamide]-tetraakis(trifluoroacetate)

A solution of the mixture of Example 16 (d) in a mixture of methylene chloride (20 mL) and trifluoroacetic acid (5 mL) was kept at r.t. for 16 hours and evaporated to dryness. A solution of the residue in a mixture of methanol (20 mL), water (2.0 mL) and 1.0 N NaOH (2.0 mL) was heated for 2 hours at 50°, cooled to r.t. and treated with trifluoroacetic acid (0.8 mL). Solvents were removed to give a pale yellow solid which was purified by semi-preparative HPLC (YMC ODS-AQ,

### Example 17

5 Preparation of N,N'-[4-(1-piperazinyl)-1,2-phenylene]bis-[1-(carboxymethyl)-4-piperidinecarboxamide]tetraakis(trifluoroacetate)

- 10 (a) N,N'-[4-(1-piperazinyl)-1,2-phenylene]bis-[1-(carboxymethyl)-4-piperidinecarboxamide]tetraakis(trifluoroacetate)

The semi-preparative hplc purification of Example 16 (c) also afforded the title compound, 0.040 g. (10.8 min.). MS (M+H)<sup>+</sup> 531.2.

### Example 18

15 Preparation of 4-[[[4-(1-Piperazinyl)phenyl]methylamino]carbonyl]-1-piperidineacetic acid

- 20 a) 4-(1-N-trifluoroacetyl-4-piperazinyl)-N-formylaniline

A solution of 4-(1-N-trifluoroacetyl-4-piperazinyl)aniline (1.2 g, 4 mmol) and Ethylformate (10 mL) was refluxed for 48 h. Reaction was concentrated. EtOAc was added and solution was washed consecutively with water, brine, dried over MgSO<sub>4</sub>, filtered and concentrated to give the title compound (0.93g, 77%). MS(ES) m/e 302 [M+H]<sup>+</sup>.

- 25 b) 4-(1-N-trifluoroacetyl-4-piperazinyl)-N-methylaniline

To a solution of compound of Example 18(a) (0.9 g, 3.1 mmol) in THF (15 mL) was added 2M BH<sub>3</sub>dimethylsulfide complex (5 mL, 10 mmol). Reaction was refluxed for 2 h. 4M HCl in Dioxane (2 mL) and MeOH (2 mL) were added and reaction was refluxed for 1 h. Reaction was concentrated and title compound was crystallized from EtOAc/MeOH (0.54g, 55%). MS(ES) m/e 361 [M+H]<sup>+</sup>.

- 30 c) Ethyl 4-chlorocarbonyl-1-piperidine acetate

A solution of ethyl 4-carboxy-1-piperidine acetate (0.44 g, 1.75 mmol) and thionyl chloride (0.2 mL, 3.9 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (3 mL) was stirred at RT for 4 h. Reaction was concentrated and carried on to the next step (0.47g, Quant.)



d) Ethyl-4-[[[4-(4-Methyl-1-piperazinyl)phenyl]methylamino]carbonyl]-1-piperidine acetate

To a stirred solution of compound of Example 18(c) (1.6 g, 1.7 mmol) in

5  $\text{CH}_2\text{Cl}_2$  (3 mL) was added a solution of compound of Example 12(b) (0.46 g, 3 mmol) and  $\text{Et}_3\text{N}$  (0.28 mL, 2 mmol) in  $\text{CH}_2\text{Cl}_2$  (3 mL). Reaction was stirred for 18 h.

Reaction was washed consecutively with water, 5%  $\text{NaHCO}_3$ , brine. Organic solution was concentrated to give the title compound (0.62g, 75%). MS(ES)m/e 485 [M+H]<sup>+</sup>.

e) 4-[[[4-(1-Piperazinyl)phenyl]methylamino]carbonyl]-1-piperidine acetic acid

A solution of Ethyl-4-[[[4-(1-Piperazinyl)phenyl]methylamino]carbonyl]-1-piperidine acetate (0.6 g, 0.23 mmol) in MeOH (3 mL) was treated with 1N NaOH (1.0 mL, 1.0 mmol). Solution was stirred at RT for 18 hr. Solution was concentrated and the residue was purified by prep HPLC (5%  $\text{CH}_3\text{CN}/0.1\%$  TFA in water) to give the title compound (0.18g, 20%). MS(ES)m/e 361 [M+H]<sup>+</sup>.

<sup>1</sup>H NMR (250MHz, DMSO-d<sub>6</sub>)  $\delta$  1.5-4.3 (m, 22H), 6.9-9.5 (m, 5H).

Anal. ( $\text{C}_{19}\text{H}_{24}\text{N}_4\text{O}_3 \cdot 3\text{TFA} \cdot 2.5\text{H}_2\text{O}$ ) calcd: C, 39.83; H, 4.00; N, 6.98. found: C, 40.03; H, 4.18; N, 7.13.

#### Example 19

Preparation of Sodium 4-[[[4-(1-piperazinyl)phenyl]carbonyl]amino]-1-piperidine acetate

a) Sodium 4-[[[4-(1-piperazinyl)phenyl]carbonyl]amino]-1-piperidine acetate

A solution of Sodium 4-[[[4-(1-benzyl-piperazinyl)phenyl]carbonyl]amino]-1-piperidine acetate (0.3 g, 0.98 mmol) and 10% Pd/C (0.1 g) in MeOH (40 mL) was hydrogenated (37 psi) for 5hr. Solution was filtered through Celite and the filtrate was concentrated to give the title compound (0.25g, 85%). MS(ES) m/e 347 [M+H]<sup>+</sup>.

<sup>1</sup>H NMR (250 MHz, DMSO-d<sub>6</sub>)  $\delta$  1.5-4.5 (m, 18H), 6.8-9.5 (m, 5H).

Anal. ( $\text{C}_{18}\text{H}_{23}\text{N}_4\text{O}_3 \cdot 0.75\text{CH}_3\text{OH}$ ) calcd: C, 57.00; H, 7.01; N, 13.98. found: C, 57.19; H, 7.23; N, 14.10.

#### Example 20

Preparation of 4-[N-[[4-(1-piperazinyl)benzyl]-N-acetyl]amino]-1-piperidine-acetic acid

a) 4-fluoro-methylbenzoate

Thionyl chloride (16mL, 0.22 mol) was added slowly to a solution of 4-fluorobenzoic acid (10 g, 71 mmol) in MeOH (200 mL). The reaction was stirred at RT for 18 h. Solution was evaporated to leave an oil (10.7 g, 97%). <sup>1</sup>H NMR (250MHz, DMSO-d<sub>6</sub>)  $\delta$  3.8 (s, 3H), 7.0-8.1 (m, 4H).

b) 4-piperazinyl-methyl benzoate

A stirred solution of piperazine (2.6 g, 30.2 mmol) and methyl ester (1.93 g, 12.1 mmol) of Example 20(a) in DMSO (15 mL) was stirred at 65°C for 18 h. Solution was cooled to RT. Reaction was treated with  $\text{NaHCO}_3$  and extracted with EtOAc. The combined extracts were concentrated to afford the title compound (2.28 g, 84%). MS(ES) m/e 221 [M+H]<sup>+</sup>.

c) 4-[1'-(N-t-butyloxycarbonyl)-piperazinyl]-methyl benzoate

Boc<sub>2</sub>O (2.5 g, 10.4 mmol) was added to a solution of the piperazine (2.28 g, 10.4 mmol) of Example 20(b) and  $\text{Et}_3\text{N}$  (1.7 mL, 11 mmol) in  $\text{CH}_3\text{CN}$  (40 mL).

The reaction was stirred at RT for 18 h. Solution was filtered and the solid was washed with  $\text{CH}_3\text{CN}$  to afford the title compound (1.88 g, 57%). MS(ES) m/e 321 [M+H]<sup>+</sup>.

d) 4-[1'-(N-t-butyloxycarbonyl)-piperazinyl]-benzoic acid

The compound of Example 20(c) (1.88 g, 5.9 mmol) was dissolved in MeOH/THF (10 mL/100 mL) and treated with 1N NaOH (12 mL, 12 mmol). The solution was stirred at RT for 18 h. The solution was concentrated and the residue was taken up in water and AcOH was added dropwise until precipitation occurred. The solution was filtered and the solid was washed with water and dried *in vacuo* to give the title compound (1.7g, 94%). MS(ES) m/e 307 [M+H]<sup>+</sup>.

e) 4-[[[4-(4-t-butyloxycarbonyl)-1-piperazinyl]phenyl]carbonyl]amino]-1-benzyl-piperidine

To a solution consisting of 4-amino-1-benzylpiperidine (1.1 mL, 4.8 mmol) in DMF (10 mL) was added  $\text{Et}_3\text{N}$  until the pH was 9. To the resulting system was added consecutively HOBt (865 mg, 4.8 mmol), the compound of Example 20(d) (1.47 g, 4.5 mmol) and water soluble carbodiimide (1.0 g, 4.8 mmol). The resulting

solution was stirred at RT for 18 h. The solution was concentrated and chromatographed using silica gel and CH<sub>2</sub>Cl<sub>2</sub>/MeOH (20/1). Desired fractions were concentrated to give the title compound. (1.84g, 80%). MS(ES) m/e 479 [M+H]<sup>+</sup>

- 5 0-4-[[[4-(4-(t-Butyloxycarbonyl)-1-piperazinyl)phenyl]carbonyl]amino]-1-piperidine

A solution consisting of the compound of Example 20(e) (1.84 g, 3.8 mmol) and Pearlman's reagent (1.4 g) in MeOH (50 mL) was hydrogenated at 50 psi for 2.5 h. Reaction was filtered through Celite and the filtrate was concentrated to afford the title compound (1.5g, 90%). MS(ES) m/e 389 [M+H]<sup>+</sup>.

- 10 g)-Butyl-4-[[[4-(4-(t-Butyloxycarbonyl)-1-piperazinyl)phenyl]carbonyl]amino]-1-piperidine acetate

A solution of compound of Example 20(f) (1.35 g, 3.5 mmol) and Et<sub>3</sub>N (0.727 mL, 4.0 mmol) in DMF (20 mL) was treated with t-butyl-bromo acetate (0.565 mL, 3.7 mmol). Reaction was stirred at RT for 18 h. Solution was concentrated to give title compound (1.65g, 94%). MS(ES) m/e 503 [M+H]<sup>+</sup>.

- 15 h) t-Butyl-4-[[[4-(4-(t-Butyloxycarbonyl)-1-piperazinyl)phenyl]carbonyl]amino]-1-piperidine acetate

A stirred mixture of compound of Example 20(g) (0.86 g, 1.7 mmol) and Lawesson reagent (0.693 g, 1.7 mmol) in toluene (20 mL) was heated to 80°C for 2 h. The solution was concentrated and chromatographed using silica gel and EtOAc/Hexane (1/1). Desired fractions were concentrated to give the title compound (0.56g, 63%). MS(ES) 519 [M+H]<sup>+</sup>.

- 20 i) t-Butyl-4-[[[4-(4-(t-Butyloxycarbonyl)-1-piperazinyl)phenyl]carbonyl]amino]-1-piperidine acetate

A stirred mixture of compound of Example 20(h) (0.86 g, 1.7 mmol) and Lawesson reagent (0.693 g, 1.7 mmol) in toluene (20 mL) was heated to 80°C for 2 h. The solution was concentrated and chromatographed using silica gel and EtOAc/Hexane (1/1). Desired fractions were concentrated to give the title compound (0.56g, 63%). MS(ES) 519 [M+H]<sup>+</sup>.

25 j) t-Butyl-4-[[[4-(4-(t-Butyloxycarbonyl)-1-piperazinyl)phenyl]carbonyl]amino]-1-piperidine acetate

Sodium borohydride (0.894 mg, 24 mmol) was added to a cooled solution of compound of Example 20(h) (0.51 g, 0.98 mmol) and NiCl<sub>2</sub>·6H<sub>2</sub>O (1.9 g, 8 mmol) in THF/MeOH (1/1). The reaction was stirred for 20 minutes. Solution was concentrated and residue was extracted with CH<sub>2</sub>Cl<sub>2</sub>. The combined extracts were concentrated to give the title compound (0.48g, 85%) MS(ES) 489 [M+H]<sup>+</sup>.

- 30 k) t-Butyl-4-[[[4-(4-(t-Butyloxycarbonyl)-1-piperazinyl)phenyl]carbonyl]amino]-1-piperidine acetate

Following the procedure of Example 20(i), the title compound was prepared (0.12g, 86%). <sup>1</sup>H NMR (250 MHz, DMSO-d<sub>6</sub>) δ 1.5-4.2 (m, 21H), 7.0-9.1 (m, 6H). MS(ES) m/e 333 [M+H]<sup>+</sup>. Anal. (C<sub>24</sub>H<sub>33</sub>N<sub>3</sub>O<sub>7</sub>·2H<sub>2</sub>O·2TFA) calcd: C, 44.30; H, 5.74; N, 9.39. found: C, 44.60; H, 5.53; N, 9.32.

Acetyl chloride (0.03 mL, 0.42 mmol) was added to a cooled solution of compound of Example 20(i) (0.205 g, 0.42 mmol) and Et<sub>3</sub>N (0.117 mL, 0.84 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (5 mL). The reaction was stirred at RT for 18 h. Solution was washed with 5% NaHCO<sub>3</sub> and H<sub>2</sub>O. Organic layer was concentrated to give the title compound (0.23g, 100%). MS(ES) 531 [M+H]<sup>+</sup>.

- 5

k) 4-[[[4-(4-(1-piperazinyl)benzyl)]-N-acetyl]amino]-1-piperidine acetic acid

A solution of compound of Example 20(j) (0.23 g, 0.43 mmol) in CH<sub>2</sub>Cl<sub>2</sub>/TFA (1/1) was stirred at RT for 18 h. Reaction was concentrated and chromatographed using C-18 reversed-phase and CH<sub>3</sub>CN (15% in water with 0.1% TFA). Desired fractions were concentrated to give the title compound (0.08g, 50%). <sup>1</sup>H NMR (250 MHz, DMSO-d<sub>6</sub>) δ 1.5-4.5 (m, 24H), 6.9-7.3 (m, 4H). MS(ES) m/e 375 [M+H]<sup>+</sup>. Anal. (C<sub>20</sub>H<sub>29</sub>N<sub>3</sub>O<sub>5</sub>·TFA·2.5 H<sub>2</sub>O) calcd: C, 49.53; H, 6.80; N, 10.50. found: C, 49.83; H, 6.83; N, 10.46.

- 10 l) 4-[[[4-(4-(1-piperazinyl)phenyl)amino]thiocarbonyl]-1-piperidine acetic acid

Following the procedure of Example 20(k) except substituting compound of Example 20(h) for compound of Example 20(j), the title compound was prepared (0.040g, 20%). <sup>1</sup>H NMR (250 MHz, DMSO-d<sub>6</sub>) δ 1.7-4.6 (m, 19H), 6.9-9.8 (m, 3H). ES(ES) m/e 363 [M+H]<sup>+</sup>. Anal. (C<sub>18</sub>H<sub>25</sub>N<sub>3</sub>O<sub>3</sub>S·1.25TFA·2H<sub>2</sub>O) calcd: C, 45.40; H, 5.78; N, 10.25. found: C, 45.48; H, 5.88; N, 10.37.

### Example 21

Preparation of 4-[[[4-(4-(1-piperazinyl)phenyl)amino]thiocarbonyl]-1-piperidine acetic acid

- 20 a) 4-[[[4-(4-(1-piperazinyl)phenyl)amino]thiocarbonyl]-1-piperidine acetic acid

Following the procedure of Example 20(k) except substituting compound of Example 20(h) for compound of Example 20(j), the title compound was prepared (0.040g, 20%). <sup>1</sup>H NMR (250 MHz, DMSO-d<sub>6</sub>) δ 1.7-4.6 (m, 19H), 6.9-9.8 (m, 3H). ES(ES) m/e 363 [M+H]<sup>+</sup>. Anal. (C<sub>18</sub>H<sub>25</sub>N<sub>3</sub>O<sub>3</sub>S·1.25TFA·2H<sub>2</sub>O) calcd: C, 45.40; H, 5.78; N, 10.25. found: C, 45.48; H, 5.88; N, 10.37.

### Example 22

- 30 Preparation of 4-[[[4-(4-(1-piperazinyl)benzyl)amino]-1-piperidine-acetic acid

a) 4-[[[4-(4-(1-piperazinyl)benzyl)amino]-1-piperidine-acetic acid

Following the procedure of Example 20(k) except substituting compound of Example 20(i) for compound of Example 20(j), the title compound was prepared (0.12g, 86%). <sup>1</sup>H NMR (250 MHz, DMSO-d<sub>6</sub>) δ 1.5-4.2 (m, 21H), 7.0-9.1 (m, 6H). MS(ES) m/e 333 [M+H]<sup>+</sup>. Anal. (C<sub>24</sub>H<sub>33</sub>N<sub>3</sub>O<sub>7</sub>·2H<sub>2</sub>O·2TFA) calcd: C, 44.30; H, 5.74; N, 9.39. found: C, 44.60; H, 5.53; N, 9.32.

**Example 23**

Preparation of 1-[[[4-(1-piperazinyl)phenyl]amino]carbonyl]-4-piperazineacetic acid

- 5 a) 4-[4-(trifluoroacetyl)-1-piperazinyl]phenylisocyanate **2**  
 4-[4-(trifluoroacetyl)-1-piperazinyl]aniline **4** (250mg, 0.9mmole) was dissolved in dioxane (5ml) and TEA (125l, 0.9mmole). Trichloromethylchloroformate (110l, 0.9mmole) was added dropwise via syringe. Solid precipitated out upon addition, mixture stirred 30 min and was filtered to yield the title compound as a white solid (100mg, 37%). MS(ES) m/e 300.2 [M+H]<sup>+</sup>
- 10 b) methyl-1-[[[4-(1-(4-trifluoroacetyl)piperazinyl]phenyl]amino]carbonyl]-4-piperazine acetate **3**  
 N-(carboethoxymethyl)piperazine **40** (280mg, 16mmole) in THF (5ml) was added dropwise to the compound of example 23 (a) dissolved in THF (10ml) at room temperature. The solution stirred overnight was concentrated and chromatographed (2-3% MeOH/CH<sub>2</sub>Cl<sub>2</sub>, silica gel) to yield the title compound as a light brown solid (1.0g, 26%). MS(ES) m/e 472.2 [M+H]<sup>+</sup>
- 15 c) 1-[[[4-(1-piperazinyl)phenyl]amino]carbonyl]-4-piperazineacetic acid tris(trifluoroacetate) salt  
 The compound of example 23 (b) (700mg, 1.5mmole) was dissolved in MeOH (10ml), 1N NaOH (4ml) and H<sub>2</sub>O (3ml) and the solution stirred overnight. The MeOH was evaporated and aqueous solution brought to pH 6.5 with 1N HOAc. This solution was concentrated and residue purified by prep HPLC (YMC ODS-AQ, 50x250mm, 10 $\mu$ m, 120A, 220nm, 80ml/min, .1% TFA-2.5% CH<sub>3</sub>CN/H<sub>2</sub>O) to yield the title compound as a white solid (37mg, 7%). MS(ES) m/e 348.2 [M+H]<sup>+</sup>

30

**Example 24**

Preparation of 1-[[[4-(1-piperazinyl)phenyl]amino]carbonyl]-4-piperidine acetic acid

- 35 a) 4-[(1-t-Butyloxycarbonyl)piperazine]-4-phenylisocyanate **42**  
 To a stirred solution of 4-[(1-t-Butyloxycarbonyl)piperazine]-4-aniline **3** (0.874 g, 3.15 mmol) in dioxane (4 mL), was added triethylamine (440  $\mu$ L, 3.15 mmol) followed by a dropwise addition of diphosgene (0.623 g, 3.15 mmol). After a

81

stirring of 2 h under argon, the triethylamine hydrochloride was filtered, and the filtrate was concentrated to afford the title compound as a white solid (0.56 g, 58.6%). MS(ESI) m/e 304.0 [M+H]<sup>+</sup>. HPLC k' 8.2 (Ultrasphere  $\Phi$  ODS, gradient, A:acetonitrile B:water-0.1% trifluoroacetic acid, 5-60% acetonitrile during 20 min; UV detection at 220 nm). IR (CH<sub>2</sub>Cl<sub>2</sub> film on KBr crystal) 2261 cm<sup>-1</sup>.

5

- b) Ethyl-4-piperidine acetate hydrochloride **44**

To a solution of ethyl-4-pyridine acetate (4.2 g, 2.5 mmol) in methanol (100 ml) was added PtO<sub>2</sub> (0.56 g, 2.5 mmol) and HCl (25 mL of 1N) under argon. The solution was hydrogenated under 50 psi for 4 h. The solution was filtered through celite, concentrated and triturated with ether to afford the title compound as a white solid (5.1 g, quantitative). MS(ESI) m/e 172 [M+H]<sup>+</sup>. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  6.25 (brs, 2H), 4.05 (q, 2H), 3.4 (m, 2H), 2.8 (m, 2H), 2.25 (d, 2H), 1.8 (d, 2H), 1.55 (d, 2H), 1.1 (t, 3H).

15

- c) Ethyl-1-[[[4-(1-t-Butyloxycarbonyl)piperazinyl]phenyl]amino]carbonyl]-4-piperidine acetate **45**

The compound of Example 24(a) (0.56 g, 1.85 mmol) was dissolved in dry THF (20 mL), treated with the compound of Example 24(b) (0.457 g, 2.2 mmol) and triethylamine (306  $\mu$ L, 2.2 mmol). The solution was stirred under argon for 24 h. The reaction was concentrated and the residue was extracted with ethyl acetate and washed with water. The organic extract was dried (anhydrous Na<sub>2</sub>SO<sub>4</sub>), filtered, concentrated and chromatographed (silica gel, ethyl acetate), then recrystallized from ethyl acetate to afford the title compound as off white solid (0.56 g, 64%). MS(ESI) m/e 475.4 [M+H]<sup>+</sup>. TLC R<sub>f</sub> 0.63 (silica gel, ethyl acetate).

25

- d) Ethyl-1-[[[4-(1-piperazinyl)phenyl]amino]carbonyl]-4-piperidine acetate

The compound of Example 24(c) (0.8 g, 1.69 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (10 ml) was added trifluoroacetic acid (10 ml) at RT and stirred for 1h. The resulting solution was then concentrated to dryness on rotavap and evaporated with CH<sub>2</sub>Cl<sub>2</sub> twice to afford the title compound. MS(ESI) m/e 375.4 [M+H]<sup>+</sup>. HPLC k' 7.18 (Ultrasphere  $\Phi$  ODS, gradient, A:acetonitrile B:water-0.1% trifluoroacetic acid, 5-60% acetonitrile during 20 min; UV detection at 220 nm).

30

- 35 e) 1-[[[4-(1-piperazinyl)phenyl]amino]carbonyl]-4-piperidine acetic acid

To a solution of the compound of Example 24(d) in ethanol (10 mL) was added a solution of 1N NaOH (12 ml) and was stirred at RT for 20 h. It was then

82

concentrated, and the resulting oily residue was dissolved in H<sub>2</sub>O, and the pH was adjusted to 7.08 by means of 50% acetic acid. The aqueous solution was purified on flash ODS column (step gradient, 4-6% acetonitrile/water. The fractions containing the pure compound were collected, concentrated and lyophilized to yield the title compound (430 mg, 73%) as a white powder. HPLC k' 5.2 (Ultrasphere ® ODS, gradient, A:acetonitrile B:water-0.1% trifluoroacetic acid, 5-60% acetonitrile during 20 min; UV detection at 220 nm). MS (ESI) m/e 347.2 [M+H]<sup>+</sup>; Anal. (C<sub>18</sub>H<sub>26</sub>N<sub>4</sub>O<sub>4</sub>, 1.5 H<sub>2</sub>O) calcd: C, 57.89; H, 7.83; N, 15.00. Found: C, 58.18; H, 7.79; N, 15.03. TLC R<sub>f</sub> 0.6 (silica gel, 1:1:1:1, ethyl acetate/butanol, water, acetic acid).

#### Example 25

Preparation of Trans-4-[[[4-(1-piperazinyl)phenyl]amino]carbonyl]-1-hydroxy-L-cyclohexanecarboxylic acid (trifluoroacetate)

a) 4-Oxo-cyclohexanecarboxylic acid 47

A mixture of 2% H<sub>2</sub>SO<sub>4</sub> and ethyl 4-oxo-cyclohexanoate 46 (10.0 g., 58.7 mmole) was stirred vigorously at 80° for 2 hours. The solution was cooled to r.t. and decanted. The supernatant was extracted with ether (3 x 150 mL.) and the combined organic layers

dried over Na<sub>2</sub>SO<sub>4</sub> and evaporated to dryness. The pale yellow residue was stirred with warm 30% ethyl acetate in hexane (200 mL.) and filtered. The title compound (3.5 g.) precipitated from the filtrate upon cooling. MS (M+H)<sup>+</sup> 143.2

b) N-[4-(4-(t-butoxycarbonyl)-1-piperazinyl)phenyl]-4-oxo-cyclohexanecarboxamide 48

A solution of the compound of Example 25 (a) (1.7 g., 12.0 mmole) in a mixture of methylene chloride (12.0 mL.) and dimethylformamide (1.9 mL.) was cooled to -40° under argon and a solution of thionyl chloride (1.43 g., 12.0 mmole) in methylene chloride (8 mL.) was added dropwise. The solution was stirred 30 minutes and a solution of 4-(4-(t-butoxycarbonyl)-1-piperazinyl)aniline 3 (3.3 g., 12.0 mmole) and triethylamine (2.4 g., 24.0 mmole) in methylene chloride (30 mL.) was added dropwise over one hour. The solution was allowed to stir at r.t. for two hours, diluted with methylene chloride (280 mL.), extracted with water (2 x 50 mL.), 5% NaHCO<sub>3</sub> (50 mL.) and water (50 mL.) and dried finally over Na<sub>2</sub>SO<sub>4</sub>. Removal

of solvent gave a yellow solid purified by flash chromatography (3% CH<sub>3</sub>OH/CH<sub>2</sub>Cl<sub>2</sub>, silica) to give the title compound, 2.0 g. MS (M+H)<sup>+</sup> 402.2

c) Ethyl trans-4-[[[4-(4-t-butoxycarbonyl)-1-piperazinyl]-phenyl]amino]carbonyl]-1-hydroxy-L-cyclohexanecarboxylate and ethyl cis-4-[[[4-(4-t-butoxycarbonyl)-1-piperazinyl]-phenyl]amino]carbonyl]-1-hydroxy-L-cyclohexanecarboxylate 49

A solution of lithium bis(trimethylsilyl)amide (1.0 M in THF, 10.0 mL., 10 mmole) was added to anhydrous THF (80 mL.). The solution was cooled to -78° and a solution of ethyl acetate (0.88 g., 10 mmole) in THF (5.0 mL.) was added dropwise over five minutes. This solution was stirred 15 minutes and a solution of the compound of Example 25 (b) (1.3 g., 3.24 mmole) in THF (65 mL.) was added dropwise over 15 minutes. The resulting mixture was stirred 25 minutes at -78°, quenched with glacial acid (1.0 mL.) and allowed to warm to r.t. Solvents were removed and the residue partitioned between ethyl acetate (300 mL.) and water (60 mL.). The organic layer was dried over Na<sub>2</sub>SO<sub>4</sub> and evaporated to dryness to give a mixture of alcohols, 1.5 g. MS (M+H)<sup>+</sup> 490.2

d) Trans-4-[[[4-(1-piperazinyl)phenyl]amino]carbonyl]-1-hydroxy-L-cyclohexanecarboxylic acid (trifluoroacetate) and cis-4-[[[4-(1-piperazinyl)phenyl]amino]carbonyl]-1-hydroxy-L-cyclohexanecarboxylic acid (trifluoroacetate)

A solution of the mixture of Example 25 (c) in a mixture of methylene chloride (30 mL.) and trifluoroacetic acid (10 mL.) was kept overnight at 0°. Solvents were removed and the residue triturated with ether to give a colorless solid, which was dissolved in a mixture of methanol (50 mL.), water (5 mL.) and 1.0 N NaOH (5.0 mL.). This solution was heated at 50° for 2 hours, cooled to r.t. and treated with trifluoroacetic acid (1.5 mL.). Solvents were removed to give a pale yellow solid which was purified by semi-preparative HPLC (YMC ODS-AQ, 10 µm, 120°, 50 X 250 mm.; 12% CH<sub>3</sub>CN/H<sub>2</sub>O, 0.1% TFA; 90 mL/min.) to give the title compound, 0.50 g. (eluting at 10.1 min.) and 0.44 g. (15.0 min.) respectively. MS (M+H)<sup>+</sup> 362.2.

#### Example 26

Preparation of cis-4-[[[4-(1-piperazinyl)phenyl]amino]carbonyl]-1-hydroxy-L-cyclohexanecarboxylic acid (trifluoroacetate)

(a) cis-4-[[[4-(1-piperazinyl)phenyl]amino]carbonyl]-1-hydroxy-1-cyclohexanecarboxylic acid (trifluoroacetate)

The semi-preparative hplc purification of Example 25 (d) also afforded the title compound, 0.44 g. (15.0 min.). MS (M+H)<sup>+</sup> 362.2.

5

#### Example 27

Preparation of 4-[[[4-(1-piperazinyl)phenyl]amino]carbonyl]-1-cyclohexanecarboxylic acid

10 a) Ethyl 4-trifluoromethanesulfonylphenyl acetate 51

A solution containing ethyl 4-hydroxyphenyl acetate 50 (4.9g, 27.2mmol) and pyridine (6.6ml, 82mmol) in CH<sub>2</sub>Cl<sub>2</sub> (sure-seal, 17 ml) was added dropwise to a solution of trifluoromethanesulfonic anhydride (6.36ml, 35.4mmol) in CH<sub>2</sub>Cl<sub>2</sub> (10ml) with cooling under argon. After 4h, the solvent was removed and the residual was triturated with Et<sub>2</sub>O-hexane, insoluble salt was removed by filtration and the filtrate was concentrated, after silica chromatography get 7.3g (86% yield) as a light amber oil: MS(ES) m/e 313 (M + H)<sup>+</sup>.

15

b) Ethyl 4-carboxyphenyl acetate 52

A mixture of the compound of Example 27 (a) (624mg, 2mmol), potassium acetate (784mg, 8mmol), palladium acetate (22.5mg, 0.1mmol) and 1,1'-Bis(diphenylphosphino)ferrocene (221.6mg, 0.4mmol) in DMSO (12ml) was purged with carbon monoxide for 10 min. and then heated under CO balloon at 60°C for 5h. The reaction mixture was diluted with water, acidified with 0.5N hydrochloric acid and extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic layer was washed with water, dried (Na<sub>2</sub>SO<sub>4</sub>), and evaporated under vacuum. The residue was chromatographed on silica gel and get 280mg (67% yield) as an off white solid: MS(ES) m/e 224 (M + NH<sub>4</sub>)<sup>+</sup>.

20

25

c) Ethyl 4-carboxycyclohexanecarboxylate 53

A mixture of the compound of 27 (b) (1.1g, 5.3mmol) and platinum(IV) oxide (602mg, 2.65mmol) in acetic acid (60ml) was hydrogenated at 50 psi in a Parr apparatus at RT for 5.5 hr. The reaction mixture was purged with argon, filtered and concentrated gave the title compound (1.01g, 89%) as a clear oil: MS (ES) m/e 215.0 (M + H)<sup>+</sup>.

30

85

d) Ethyl 4-[[[4-(1-(tert-butoxycarbonyl)piperazinyl)phenyl]amino]carbonyl]-1-cyclohexanecarboxylate 54

Following the procedure of Example 3 (a), except substituting the compound of Example 27 (c) for Ethyl 4-carboxypiperidinyl acetate 8, the title compound (1.9g, 85.5%) as a pale yellow solid: (ES) m/e 474 (M + H)<sup>+</sup>.

5

e) Ethyl 4-[[[4-(1-piperazinyl)phenyl]amino]carbonyl]-1-cyclohexanecarboxylate:

Following the procedure of Example 3 (b), except substituting the compound of Example 27 (d) for Ethyl 4-[[[4-(1-(tert-butoxycarbonyl)piperazinyl)phenyl]amino]carbonyl]-1-piperidine acetate 16, the title compound (71%) as a white solid: (ES) m/e 374 (M + H)<sup>+</sup>.

10

f) 4-[[[4-(1-piperazinyl)phenyl]amino]carbonyl]-1-cyclohexanecarboxylic acid

Following the procedure of Example 8 (c), the compound of Example 27 (e) was saponified gave the title compound as a white powder: MS (ES) m/e 346.2 (M + H)<sup>+</sup>. Anal. Calcd for C<sub>19</sub>H<sub>27</sub>N<sub>3</sub>O<sub>3</sub> 1 HCl 6 0.33 H<sub>2</sub>O: C, 58.83; H, 7.45; N, 10.83. Found: C, 59.10; H, 7.56; N, 10.81.

15

#### Example 28

Preparation of 4-[[[4-(1-piperazinyl)phenyl]amino]carbonyl]-1-piperidine propionic acid

20

a) 4-(1-N-trifluoroacetyl-4-piperazinyl)nitrophenol

Trifluoroacetic anhydride (7.5 mL, 48.5 mmol) was added to a cooled solution of 4-piperazinylnitrophenol (10 g, 48.3 mmol) and Et<sub>3</sub>N (13.5 mL, 96 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (100 mL). Reaction was stirred at RT for 1 h. Solution was washed consecutively with 5% NaHCO<sub>3</sub>, H<sub>2</sub>O, 5% citric acid and H<sub>2</sub>O. The organic layer was concentrated. The crude was purified by flash chromatography (silica gel, EtOAc/Hexane (1/1)) to yield the title compound (12.1g, 83%). MS(ES) m/e 304 [M+H]<sup>+</sup>.

25

30

b) 4-(1-N-trifluoroacetyl-4-piperazinyl)aniline

To a solution of 4-(1-N-trifluoroacetyl-4-piperazinyl)nitrophenol (2.0 g, 6.6 mmol) in DMF/MeOH (20 mL/10 mL) was added 10% Pd/C (0.8 g). Solution was hydrogenated (45 psi) for 2 hr. Solution was filtered through Celite and the filtrate was concentrated to give the title compound (1.9g, 100%). MS(ES) m/e 274 [M+H]<sup>+</sup>.

35

86

a) 4-[[[4-(1-N-trifluoroacetyl-4-piperazinyl)phenyl]amino]carbonyl]-1-N-1-butylloxycarbonylpiperidine

A mixture of 4-(1-N-trifluoroacetyl-4-piperazinyl)aniline (3.0 g, 9.9 mmol), Et<sub>3</sub>N (2.2 mL, 19 mmol), HOBT (1.5 g, 9.9 mmol), 1-t-butylloxycarbonyl-4-carboxypiperidine (2.3 g, 9.9 mmol) and EDC (1.9 g, 9.9 mmol) in DMF (15 mL) was stirred at RT for 18 hr. Solution was concentrated and the residue was dissolved in EtOAc. EtOAc was washed with: H<sub>2</sub>O, 5% citric acid (2X), water, 5% NaHCO<sub>3</sub> (2X), water and brine. EtOAc was concentrated to give the title compound (4.35g, 91%). MS(ES) m/e 485 [M+H]<sup>+</sup>.

b) 4-[[[4-(1-N-trifluoroacetyl-4-piperazinyl)phenyl]amino]carbonyl]-1-piperidine

4-[[[4-(1-N-trifluoroacetyl-4-piperazinyl)phenyl]amino]carbonyl]-1-N-1-butylloxycarbonylpiperidine (2.0 g, 4.1 mmol) was treated with 4M HCl (30 mL) for 1 hr. Et<sub>2</sub>O (100 mL) was added. Filtration gave the title compound (1.97g, 100%). MS(ES) m/e 385 [M+H]<sup>+</sup>.

c) Methyl-4-[[[4-(1-N-trifluoroacetyl-4-piperazinyl)phenyl]amino]carbonyl]-1-piperidine acetate

To a solution of 4-[[[4-(1-N-trifluoroacetyl-4-

piperazinyl)phenyl]amino]carbonyl]-1-piperidine (0.90 g, 2.1 mmol) and Et<sub>3</sub>N (0.75 mL, 4.5 mmol) in DMF (5 mL) was added Methylbromoacetate (0.165 mL, 2.3 mmol). Solution was stirred at RT for 18 hr. Solution was concentrated and residue was taken up in EtOAc. EtOAc was washed successively with 5% citric acid (2X), water, 5% NaHCO<sub>3</sub> (2X) and water. EtOAc was concentrated to give the title compound (0.200g, 20%). MS(ES) m/e 473 [M+H]<sup>+</sup>.

d) 4-[[[4-(1-Piperazinyl)phenyl]amino]carbonyl]-1-piperidine propionic acid

Methyl-4-[[[4-(1-N-trifluoroacetyl-4-piperazinyl)phenyl]amino]carbonyl]-1-piperidine acetate (1.1 g, 2.1 mmol) was treated with 1N NaOH (4.5 mL, 4.5 mmol). Solution was stirred at RT for 18 hr. Solution was concentrated and chromatographed (C-18, 4% CH<sub>3</sub>CN, 0.1% H<sub>2</sub>O) to give the title compound (0.294g, 39%). <sup>1</sup>H NMR (250MHz, DMSO-d<sub>6</sub>) δ 1.5-4.0 (m, 20H), 6.8-9.7 (m, 5H). MS(ES)

### Example 29

#### Parenteral Dosage Unit Composition

A preparation which contains 20 mg of the compound of Example 1 as a sterile dry powder is prepared as follows: 20 mg of the compound is dissolved in 15 mL of distilled water. The solution is filtered under sterile conditions into a 25 mL multi-dose ampoule and lyophilized. The powder is reconstituted by addition of 20 mL of 5% dextrose in water (D5W) for intravenous or intramuscular injection. The dosage is thereby determined by the injection volume. Subsequent dilution may be made by addition of a metered volume of this dosage unit to another volume of DSW for injection, or a metered dose may be added to another mechanism for dispensing the drug, as in a bottle or bag for IV drip infusion or other injection-infusion system.

### Example 30

#### Oral Dosage Unit Composition

A capsule for oral administration is prepared by mixing and milling 50 mg of the compound of Example 1 with 75 mg of lactose and 5 mg of magnesium stearate. The resulting powder is screened and filled into a hard gelatin capsule.

### Example 31


#### Oral Dosage Unit Composition

A tablet for oral administration is prepared by mixing and granulating 20 mg of sucrose, 150 mg of calcium sulfate dihydrate and 50 mg of the compound of Example 1 with a 10% gelatin solution. The wet granules are screened, dried, mixed with 10 mg starch, 5 mg talc and 3 mg stearic acid; and compressed into a tablet.

The foregoing is illustrative of the making and using of this invention. This invention, however, is not limited to the precise embodiments described herein, but encompasses all modifications within the scope of the claims which follow.

What is claimed is:

1. A compound of the formula:




wherein:

R<sup>1</sup> is hydrogen, C<sub>1-6</sub>alkyl, HC≡C-CH<sub>2</sub>, F<sub>3</sub>C-CH<sub>2</sub>,



10 D is



A<sup>1</sup>, A<sup>2</sup>, A<sup>3</sup>, A<sup>4</sup>, and A<sup>5</sup> independently are N or CH, with the proviso that A<sup>1</sup> and A<sup>4</sup> are not simultaneously N;

Y and Y<sup>1</sup> independently are hydrogen, C<sub>1-6</sub>alkyl, halo, CF<sub>3</sub>,

15 -(CH<sub>2</sub>)<sub>n</sub>-phenyl or together form a phenyl ring;

X is C(O)NR<sup>2</sup>, NR<sup>2</sup>C(O), SO<sub>2</sub>NR<sup>2</sup>, NR<sup>2</sup>SO<sub>2</sub>, C(O), C(R<sup>2</sup>)<sub>2</sub>, S, O, NR<sup>2</sup>, CR<sup>2</sup>-OR<sup>2</sup>, C(O)C(R<sup>2</sup>)<sub>2</sub>, C(R<sup>2</sup>)<sub>2</sub>C(O), CH<sub>2</sub>CH<sub>2</sub>, CHOCH<sub>2</sub>, CH<sub>2</sub>CHOH, or CR<sup>2</sup>=CR<sup>2</sup>;

Q is a single bond or C(O);


20 Z is hydrogen, C<sub>1-6</sub>alkyl, OR<sup>2</sup>, CH<sub>2</sub>OR<sup>2</sup>, CH<sub>2</sub>CO<sub>2</sub>R<sup>2</sup>, CH<sub>2</sub>CONR<sup>2</sup>R<sup>2</sup>, CO<sub>2</sub>R<sup>2</sup>, CONR<sup>2</sup>R<sup>2</sup>, CH=CH-, -C≡C-, COR<sup>2</sup>, or -(CH<sub>2</sub>)<sub>n</sub>-phenyl;

each n independently is 0-3; and

each R<sup>2</sup> independently is H or C<sub>1-6</sub>alkyl;

or a pharmaceutically acceptable salt thereof.

2. A compound according to claim 1 of the formula:



wherein:

Y and Y<sup>1</sup> independently are hydrogen, C<sub>1-6</sub>alkyl, halo, CF<sub>3</sub>, or together form a phenyl ring;

X is C(O)NR<sup>2</sup>, NR<sup>2</sup>C(O), SO<sub>2</sub>NR<sup>2</sup>, NR<sup>2</sup>SO<sub>2</sub>, C(O), C(R<sup>2</sup>)<sub>2</sub>, S, O, or NR<sup>2</sup>;

5 Z is hydrogen, C<sub>1-4</sub>alkyl, OR<sup>2</sup>, CH<sub>2</sub>OR<sup>2</sup>, CH<sub>2</sub>CO<sub>2</sub>R<sup>2</sup>, or CO<sub>2</sub>R<sup>2</sup>; and

n is 0 or 1.

3. A compound according to claim 2 wherein A<sup>1</sup> is N, A<sup>2</sup> is CH, X is C(O)NR<sup>2</sup> or NR<sup>2</sup>C(O), Z is hydrogen, R<sup>1</sup> is hydrogen or C<sub>1-4</sub>alkyl, and R<sup>2</sup> is hydrogen or C<sub>1-4</sub>alkyl.

4. A compound according to claim 3 which is 4-[[[4-(1-piperazinyl)-phenyl]amino]carbonyl]-1-piperidineacetic acid or a pharmaceutically acceptable salt thereof.

5. A pharmaceutical composition comprising a compound according to claim 1 and a pharmaceutically acceptable carrier.

6. A method for effecting inhibition of platelet aggregation which comprises administering a compound according to claim 1.

7. A method for treating stroke or a transient ischemia attack or myocardial infarction which comprises administering a compound according to claim 1.

8. A method for promoting reperfusion of an artery or vein and inhibiting reocclusion which comprises administering a fibrinolytic agent and a compound according to claim 1.

INTERNATIONAL SEARCH REPORT		International application No. PCT/US97/00326
<b>A. CLASSIFICATION OF SUBJECT MATTER</b> IPC(6) : C07D 401/12, 403/12, 293/133; A61K 31/495, 31/445. US CL : Please See Extra Sheet. According to International Patent Classification (IPC) or to both national classification and IPC		
<b>B. FIELDS SEARCHED</b> Minimum documentation searched (classification system followed by classification symbols) U.S. : 544337,360,364,365,393; 546/186,189; 514/232, 255,315,316,317,329,330. Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) CAS ONLINE STRUCTURE SEARCH FILE BELSTEIN STRUCTURE SEARCH		
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X,P	Chemical Abstracts, Vol. 125, No. 15, 07 October 1996 (Columbus Ohio, USA), page 1216 column 2, the abstract no. 195688, PIEPER, H. "Preparation of 1-(piperazinocarbonyl)piperidine-4-alkanoates and analogs as cell aggregation inhibitors." PCT Int. Appl. WO 96 20,173, 04 July 1996 (Gerl).	1,5-8
X,P	DE 44 48 301 A1 (H. PIEPER) 27 June 1996. See entire document especially pages 20-29.	1,5-8
X	DE 42 41 632 A1 (F. HIMMELSBACH) 16 June 1994, see entire document especially pages 13-19.	1,5-8
---		2
Y		
* Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "B" earlier document published on or after the international filing date of the invention but not taken into account in the prior art search "C" document which may have priority claims or which is cited to establish the publication date of another claim or other special reasons (as specified) "D" document referring to an oral disclosure, use, exhibition or other means "E" document published prior to the international filing date but later than the priority date claimed "F" document published after the international filing date or priority date but not taken into account in the prior art search		
Date of the actual completion of the international search 13 MARCH 1997		Date of mailing of the international search report 07 APR 1997
Name and mailing address of the ISA/US Box PCT Washington, D.C. 20531		Authorized officer EMILY BERNHARDT
Facsimile No. (703) 305-3230		Telephone No. (703) 308-1235

Form PCT/ISA/210 (second sheet) (July 1992)\*

INTERNATIONAL SEARCH REPORT		International application No. PCT/US97/00326
<b>Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)</b> This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons: 1. <input type="checkbox"/> Claims Nos.: because they relate to subject matter not required to be searched by this Authority, namely:		
2. <input type="checkbox"/> Claims Nos.: because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:		
3. <input type="checkbox"/> Claims Nos.: because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).		
<b>Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)</b> This international Searching Authority found multiple inventions in this international application, as follows: Please See Extra Sheet.		
1. <input type="checkbox"/> As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.		
2. <input checked="" type="checkbox"/> As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.		
3. <input type="checkbox"/> As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:		
4. <input type="checkbox"/> No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:		
Remark on Protest <input type="checkbox"/>		The additional search fees were accompanied by the applicant's protest. <input type="checkbox"/> No protest accompanied the payment of additional search fees. <input type="checkbox"/>

Form PCT/ISA/210 (continuation of first sheet) (July 1992)\*



# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US97/00326

## A. CLASSIFICATION OF SUBJECT MATTER:

US CL :

544/357,360,364,365,393; 546/186,189; 514/252, 255,315,316,317,329,330.

## BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

Group I claims 1-8 drawn to compounds, compositions and use where  $A' = N$ .  
Group II claims 1-2,5-8 drawn to compounds, compositions and use where  $A' = CH$ .

Groups I-II are not so linked as to form a single general inventive concept as required by PCT Rule 13.1. The groups relate to compounds of considerable structural dissimilarity which as a result are separately classified and are made and used independently of each other and are not art-recognized equivalents.

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**